

# **VEGETABLE AND SMALL FRUIT CROPS - TREE NUTS**

## General Information

### GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

### Preventive Use

For cutworm, armyworm grape colapsis, or stalk borer control, STEED Insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control.

### Limitations, Restrictions, and Exceptions

#### VEGETABLE AND SMALL FRUIT CROPS

#### TREE NUTS

#### Restrictions

- PHI: Do not apply within 7 days of harvest for all crops except for pecan. Do not apply within 21 days of harvest for pecans.
- Application Interval: Do not make applications less than 15 days apart.
- Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.

- Grazing: Do not graze livestock in treated orchards or cut treated cover crops for feed.

## Remarks

Application by ground:

Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.

Application by air:

Apply the specified dosage in a minimum of 10 gallons of finished spray per acre.

Aphid: Coverage is essential for control of this pest. Under heavy outbreaks tank mixing with another product that is labeled for this pest is recommended for control.

## Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

## Rates

[field rates 0](#)

[field rates 1](#)

•

## Restricted Entry Interval

12 hours

## Timings

[N.A.](#)