

GINSENG - PHYTOPHTHORA ROOT ROT

General Information

GENERAL INFORMATION

Ridomil Gold GR is a systemic fungicide which provides control of certain diseases caused by members of the Oomycete class of fungi on selected crops.

Note: Ridomil Gold GR is a systemic fungicide having a specific mode of action and could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from insensitive strains of fungi. If treatment is not effective following the use of Ridomil Gold GR as recommended, an insensitive resistant strain of fungi may be present. If the treatment is ineffective due to the presence of a Ridomil Gold GR insensitive strain of fungi, neither Ridomil Gold GR nor any other fungicide with similar action will effectively control that disease. Consideration should then be given to the prompt use of other types of suitable fungicides. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance in your particular crop and disease control situation.

Do not use in nurseries, turf, or landscape plantings, unless specified in the label.

THE LABEL IS FOR FIELD USE ONLY AND IS NOT INTENDED FOR USE ON TRANSPLANT TRAYS, GREENHOUSES, LATH HOUSES, FLOAT HOUSES, HYDROPONIC PRODUCTION, OR IN BEDDING PLANT STRUCTURES.

Please refer to the label for the formula in calculating the amount of Ridomil Gold GR needed for band treatment.

Preharvest Intervals

Unless otherwise stated in the directions for use for a specific crop, there is no preharvest interval for Ridomil Gold GR.

Limitations, Restrictions, and Exceptions

GINSENG

Phytophthora root rot in Ginseng.

Ridomil Gold GR applied to the soil before early spring growth followed by additional applications at monthly intervals will control Phytophthora root rot in ginseng caused by Phytophthora cactorum.

Apply Ridomil Gold GR at 15 lbs./A uniformly to the soil surface in the spring before the plants begin growing.

Make up to 4 additional applications of Ridomil Gold GR at monthly intervals at 10 lbs./A.

Notes: To avoid possible illegal residues: (1) Do not apply more than a total of 60 lbs. of Ridomil Gold GR per acre of ginseng per growing season. (2) If Ridomil Gold SL is applied to the soil before early spring growth, do not apply more than 45 lbs. of Ridomil Gold GR as supplemental applications. (3) Do not harvest ginseng within 9 days after a Ridomil Gold GR application.

Method

[Surface](#)

Pre-Harvest Interval

9 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

48 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[In the spring before the plants begin growing.](#)