

TURFGRASS - CRANEFLIES

General Information

APPLICATION TO TURFGRASS

Sites of Application

Meridian 0.33G can be used for applications to turfgrasses on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, athletic fields and sod farms.

Application Methods

Apply Meridian 0.33G as a broadcast application to turfgrass for control of targeted pests. With calibrated granular equipment, uniformly broadcast Meridian 0.33G over the area being treated. Maintain adequate soil moisture before and after application for optimum control and healthy turfgrass growth. Excessively wet or dry conditions may impact the performance of Meridian 0.33G against white grubs and mole crickets. Do not mow the turf until the treated area has been irrigated or rainfall has occurred to allow for maximum and uniform uptake into the turfgrass or soil.

Application Timing

Meridian 0.33G is effective when applied at labeled rates to help prevent and control white grubs. To help prevent white grubs, apply Meridian 0.33G up to 45 days before the historical peak of adult flight for the targeted species. Irrigate within seven (7) days of application to move Meridian 0.33G into the root zone where grubs feed. To control white grubs apply Meridian 0.33G from egg hatch to 2nd instar (grubs less than half their full size). For optimum white grub control, apply Meridian 0.33G from peak adult flight through peak egg hatch of the primary species being targeted. Irrigate within one (1) day of application to move Meridian 0.33G into the root zone where grubs feed.

Limitations, Restrictions, and Exceptions

APPLICATION TO TURFGRASS

Sites of Application

Meridian 0.33G can be used for applications to turfgrasses on golf courses, residential lawns, commercial grounds (i.e., office and shopping complexes, airports), parks, playgrounds, athletic fields and sod farms.

Application Methods

Apply Meridian 0.33G as a broadcast application to turfgrass for control of targeted pests. With calibrated granular equipment, uniformly broadcast Meridian 0.33G over the area being treated. Maintain adequate soil moisture before and after application for optimum control and healthy turfgrass growth. Excessively wet or dry conditions may impact the performance of Meridian 0.33G against white grubs and mole crickets. Do not mow the turf until the treated area has been irrigated or rainfall has occurred to allow for maximum and uniform uptake into the turfgrass or soil.

Application Timing

Meridian 0.33G is effective when applied at labeled rates to help prevent and control white grubs. To help prevent white grubs, apply Meridian 0.33G up to 45 days before the historical peak of adult flight for the targeted species. Irrigate within seven (7) days of application to move Meridian 0.33G into the root zone where grubs feed. To control white grubs apply Meridian 0.33G from egg hatch to 2nd instar (grubs less than half their full size). For optimum white grub control, apply Meridian 0.33G from peak adult flight through peak egg hatch of the primary species being targeted. Irrigate within one (1) day of application to move Meridian 0.33G into the root zone where grubs feed.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

-

Restricted Entry Interval

12 hours

Timings

At oviposition (egg lay).