

ON ALMONDS, PEANUTS, PISTACHIOS AND WALNUTS IN BULK OR IN BAGS

General Information

GENERAL USE DIRECTIONS

IMPORTANT: It is recommended that the final spray mix be buffered to a pH of 5.5-7.0. Outside of this range pyrethrins can degrade and the product will lose effectiveness.

To avoid possible harm to honey bees and other beneficial insects, it is advisable to apply when the non-target insects are least active.

Pyrethrins degrade rapidly in sunlight. To ensure the best performance, coverage is key. Use spray equipment that will provide the best coverage and direct contact with as many insects as possible.

GENERAL USE DIRECTIONS FOR INDOOR APPLICATIONS

Remove pets and birds and cover fish aquariums before spraying.

Kills only at time of application. Listed insect pests must be directly contacted to be effective.

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm. Patients should be removed prior to treatment. If application is done as a surface spray, patients may re-enter once the spray has dried. If application is done as a space spray, the rooms should be ventilated for two hours after spraying. Do not return patients to room until after ventilation.

Do not apply in institutions (including libraries, schools, sports facilities, etc.) in the immediate area where occupants are present. Do not apply to classrooms while in use. Vacate rooms prior to treatment. If application is done as a surface spray, occupants may re-enter once the spray has dried. If application is done as a space spray, rooms should be ventilated for two hours after spraying. Do not allow occupants to enter room until after ventilation.

FOOD AREA USE: Avoid contamination of food or feedstuffs. Do not apply directly to

food. Do not apply space spray while food processing is underway. Foods should be removed or covered during treatment. All food processing surfaces and equipment should be covered or washed with an effective cleaning compound followed by a potable water rinse. In the home all food processing surfaces, dishes and utensils should be covered during treatment or thoroughly washed before use.

Limitations, Restrictions, and Exceptions

ON ALMONDS, PEANUTS, PISTACHIOS AND WALNUTS IN BULK OR IN BAGS:

To kill the following stored product insects: Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cigarette Beetles, Confused Flour Beetles, Drugstore Beetles, Flat Grain Beetles, Granary Weevils, Indian Meal Moths, Lesser Grain Borers, Maize Weevils, Mediterranean Flour Moths, Merchant Grain Beetles, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Saw-toothed Grain Beetles and Square-necked Grain Beetles, dilute 2.0 fluid ounces of PyGanic® PRO per gallon of water and apply as a coarse wet spray over the top of stored nuts or the outside surface of stacked bagged nuts at the rate of 4 gallons per 1,000 square feet. Apply at weekly intervals for about 6 weeks and then at 15-day intervals. The first two applications should be applied at the rate of 4 gallons per 1,000 square feet and subsequent treatments should be applied at the rate of 2 gallons per 1,000 square feet.

- NOT REGISTERED FOR THIS USE IN THE STATE OF CALIFORNIA

Method

[Spray](#)

Rates

[field rates 0](#)

•

Timings

[N.A.](#)