

INDOOR USE—DILUTED - SPACE SPRAY

General Information

DIRECTIONS FOR USE

Do not apply space spray while food processing is underway. Cover or remove all food processing surfaces. Thoroughly wash all food processing surfaces before reuse. After spraying in bakeries, meat packing plants, food processing plants, etc., all benches, shelving, equipment, etc. where exposed food will be handled must be washed with an effective cleaning compound followed by a potable water rinse to remove all traces of contamination. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above in those establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs.

Do not apply directly to food, water, or feed supplements. Do not contaminate milk or milk handling equipment. Wash udder and teats of dairy animals before milking.

Limitations, Restrictions, and Exceptions

INDOOR USE—DILUTED

SPACE SPRAY: For use in mechanical fogging or spraying equipment to control accessible stages of Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Mediterranean Flour Moths, Mosquitoes, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Yellow Mealworms, Dried Fruit Beetles, Warehouse Beetles, Rusty Grain Beetles, Lesser Grain Beetles, Coffee Bean Weevils and Khapra Beetles. Remove or cover exposed food and cover food-handling surfaces in edible product areas of food handling establishments. Close room and shut off all air conditioning or ventilating equipment. Dilute at the rate of 1 part concentrate into 1 to 10 parts water and mix well. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space, filling the room with mist. Vacate treated area and ventilate before re-entry.

Refer in the label for the Dilution table for ULV foggers.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[N.A.](#)