

## **INDOOR USE—DILUTED - TREATMENT OF STORED GRAIN**

### General Information

#### DIRECTIONS FOR USE

Do not apply space spray while food processing is underway. Cover or remove all food processing surfaces. Thoroughly wash all food processing surfaces before reuse. After spraying in bakeries, meat packing plants, food processing plants, etc., all benches, shelving, equipment, etc. where exposed food will be handled must be washed with an effective cleaning compound followed by a potable water rinse to remove all traces of contamination. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above in those establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs.

Do not apply directly to food, water, or feed supplements. Do not contaminate milk or milk handling equipment. Wash udder and teats of dairy animals before milking.

#### Limitations, Restrictions, and Exceptions

#### INDOOR USE—DILUTED

**TREATMENT OF STORED GRAIN:** To control accessible stages of: Indian Meal Moths and Angoumois Grain Moths: Apply diluted spray to the surface of binned grain at a rate of 0.5 gallons per 1,000 cubic feet. Apply at monthly intervals with the first appearance of infestation. In severe infestations of flying moths, break up webbing with a rake before spraying and make a second application in two weeks.

#### Method

[Spray](#)

#### Rates

[field rates 0](#)

- 

#### Timings

[At monthly intervals with the first appearance of infestation.](#)