

LAWN AND PERIMETER - ARMYWORMS ETC.

General Information

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

DO NOT apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 0.4 lb. a.i. per acre per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

DO NOT broadcast apply more than 0.4 lbs. a.i. (200 lbs. of Bifenthrin L/P .2 G) per Acre per year.

Not for use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

Resistance:

When products are used repeatedly for control, some insects are known to build up resistance to products used, but resistance can not be predicted. This product should conform to resistance management strategies established for the use area. Check with your local or state pest management authorities for more information.

This product, or other products with comparable modes of action, may not provide sufficient control if resistance should develop in your area. A resistant species may be present if poor efficacy can not be linked to extreme weather conditions or improper treatment. Consult pest management advisors for the other methods of control for your area if you believe resistance is a factor.

LAWN & PERIMETER APPLICATIONS

Use appropriate application equipment to broadcast Bifenthrin L/P .2 G to achieve thorough coverage of the area being treated.

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Bifenthrin L/P .2 G if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Limitations, Restrictions, and Exceptions

Armyworms, Cutworms, and Sod Webworms: For best results water the treated area with up to 0.1 inches of water to activate (release from the granule) the insecticide.

Perimeter Treatment: Treat the ground area 5 to 10 feet wide around and adjacent to the structure.

Boundary Treatment: Treat mulch areas, soil areas, vegetation areas, ornamental gardens, planter areas, pavement crevices and fence lines.

Method

[Broadcast](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

•

Timings

[N.A.](#)