

COLLARD

General Information

RECOMMENDATIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. Apply the following recommended rates in sufficient water to thoroughly cover one acre. By ground, apply a minimum of 200 gallons per acre on stone fruits and nuts; a minimum of 300 gallons per acre on pome fruits; a minimum of 200 gallons per acre on bush and vine fruits and a minimum of 10 gallons per acre on vegetable and row crops. By air, a minimum of 2 gallons per acre on small fruits and tree crops. Use the designated amount in 100 gallons of water unless otherwise specified.

Limitations, Restrictions, and Exceptions

COLLARDS

Harlequin Cabbage Bug—Use 1 pint per 100 gallons per acre. Leafhoppers, Leaf Miners—Use $\frac{1}{2}$ pint per acre. Aphids Use— $1\frac{1}{2}$ to 2 pints per 100 gallons.

Caterpillars—Use 2 pints per acre. Cabbage Looper, Diamond Back Moth, Imported Cabbage Worm—Use 2.8 pints per acre and combine with other recommended insecticide.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When pests first appear.](#)