

# CORN SWEET

## General Information

### RECOMMENDATIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. Apply the following recommended rates in sufficient water to thoroughly cover one acre. By ground, apply a minimum of 200 gallons per acre on stone fruits and nuts; a minimum of 300 gallons per acre on pome fruits; a minimum of 200 gallons per acre on bush and vine fruits and a minimum of 10 gallons per acre on vegetable and row crops. By air, a minimum of 2 gallons per acre on small fruits and tree crops. Use the designated amount in 100 gallons of water unless otherwise specified.

### Limitations, Restrictions, and Exceptions

#### CORN

Sap Beetle—Use 1½ pints per acre. Begin treatment when 10% of ears show silk. Repeat at 3 to 5 day intervals until 4 to 5 applications are made. Maize for grain or forage: Young Grasshoppers—Use 1½ pints per acre. Apply when nymphs are young. Injury may occur in the whorl and silk stage, using this type Malathion product.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Pre-Harvest Interval

5 days

#### Rates

[field rates 0](#)

•

#### Restricted Entry Interval

12 hours

#### Timings

When nymphs are young.