

RICE

General Information

RECOMMENDATIONS FOR USE

Apply when pests first appear and repeat as necessary to maintain control. Apply the following recommended rates in sufficient water to thoroughly cover one acre. By ground, apply a minimum of 200 gallons per acre on stone fruits and nuts; a minimum of 300 gallons per acre on pome fruits; a minimum of 200 gallons per acre on bush and vine fruits and a minimum of 10 gallons per acre on vegetable and row crops. By air, a minimum of 2 gallons per acre on small fruits and tree crops. Use the designated amount in 100 gallons of water unless otherwise specified.

Limitations, Restrictions, and Exceptions

RICE

Rice Leafminer—Use 2½ pints per acre. Apply when the eggs and larvae are abundant on the seedling rice and repeat as necessary. Rice Stink Bug— Use 1 to 1½ pints per acre. Apply during the early milk and dough stage of growing rice. The rice herbicide, Propanil, should not be applied within 15 days of a Malathion treatment. Broadcast use only over intermittently flooded areas. Application may not be made around bodies of water where fish or shellfish are grown and/or harvested commercially.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

Rice Leafminer: When the eggs and larvae are abundant on the seedling rice.

Rice Stink Bug: During the early milk and dough stage of growing rice.