

# **FLAX**

## General Information

### DIRECTIONS FOR USE

**ADDITIONAL REQUIREMENTS FOR AERIAL APPLICATIONS:** The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

**ADDITIONAL REQUIREMENTS FOR GROUND BOOM APPLICATION:** Do not apply with a nozzle height greater than 4 feet above the crop canopy.

**EFFECTIVENESS ON WEEDS:** This product is effective on a large number of broadleaf weeds. On Mustard, Thistle, and White top or Hoary cress, it has superiority over 2,4-D. The following are some of the weeds killed: Beggartick, Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Catsear, Chervil, Cocklebur, Croton, Daisy, Dandelion, Dock, Dragonhead mint, Dwarf nettle, Fanweed, Fat hen, Fennel, Field bindweed, Galinsoga, Goatsbeard, Goldenrod, Halbertleaved salt brush, Hempnettle, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarter (Goosefoot), Marshelder, Mexican poppy, Miner's lettuce, Narrow leaf plantain, Pennycress, Pepperweed, Perennial morninglory, Perennial ragweed, Pigweed, Poison hemlock, Prickly lettuce, Puncturevine, Purslane, Red root, Redstem, Shepherdspurse, Sicklepod, Small plantain, Sneezeweed, Sowthistle, Spanishneedle, Stinging nettle, Stinkweed, Sunflower, Thornapple, Tree-of-heaven, Vetch, Whitebrush, Wild carrot, Wild gooseberry, Wild jute, Wild marigold, Wild petunia, Wild radish, Wild sage, Witchweed, Yellow charlock-mustard, Yellow daisy, Yellow rocket.

### SELECTIVE SPRAYING

**NOTE:** When using on grains and flax - do not forage or graze dairy and meat

animals on treated areas within 7 days of slaughter. Also, except for small underseeded grains, use at least 8.0 to 10.0 gallons of water per acre for ground application and at least 1.0 to 5.0 gallons of water per acre for aerial application.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixing with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of the material not in strict accordance with directions given herewith.

#### Limitations, Restrictions, and Exceptions

FLAX: Use 3.0 to 6.0 ounces per acre. Use lower rate where susceptible weeds such as Mustard are the main problem. Use higher rate when weeds such as Pigweed, Lambsquarters, Stinkweed, Ragweed, and Cocklebur are the problem. Apply only when weeds are up and when flax is 2 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada thistle is present, it may be necessary to go as high as 0.375 pint per acre to prevent seed head production. Some injury to flax may result.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Rates

[field rates 0](#)

- 

#### Restricted Entry Interval

12 hours

#### Timings

[When weeds are up and when flax is 2 to 8 inches high and before it comes into bud.](#)