

NON-CROP SPRAYING

General Information

DIRECTIONS FOR USE

ADDITIONAL REQUIREMENTS FOR AERIAL APPLICATIONS: The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

ADDITIONAL REQUIREMENTS FOR GROUND BOOM APPLICATION: Do not apply with a nozzle height greater than 4 feet above the crop canopy.

EFFECTIVENESS ON WEEDS: This product is effective on a large number of broadleaf weeds. On Mustard, Thistle, and White top or Hoary cress, it has superiority over 2,4-D. The following are some of the weeds killed: Beggartick, Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Catsear, Chervil, Cocklebur, Croton, Daisy, Dandelion, Dock, Dragonhead mint, Dwarf nettle, Fanweed, Fat hen, Fennel, Field bindweed, Galinsoga, Goatsbeard, Goldenrod, Halbertleaved salt brush, Hempnettle, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarter (Goosefoot), Marshelder, Mexican poppy, Miner's lettuce, Narrow leaf plantain, Pennycress, Pepperweed, Perennial morninglory, Perennial ragweed, Pigweed, Poison hemlock, Prickly lettuce, Puncturevine, Purslane, Red root, Redstem, Shepherdspurse, Sicklepod, Small plantain, Sneezeweed, Sowthistle, Spanishneedle, Stinging nettle, Stinkweed, Sunflower, Thornapple, Tree-of-heaven, Vetch, Whitebrush, Wild carrot, Wild gooseberry, Wild jute, Wild marigold, Wild petunia, Wild radish, Wild sage, Witchweed, Yellow charlock-mustard, Yellow daisy, Yellow rocket.

SELECTIVE SPRAYING

NOTE: When using on grains and flax - do not forage or graze dairy and meat

animals on treated areas within 7 days of slaughter. Also, except for small underseeded grains, use at least 8.0 to 10.0 gallons of water per acre for ground application and at least 1.0 to 5.0 gallons of water per acre for aerial application.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixing with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of the material not in strict accordance with directions given herewith.

Limitations, Restrictions, and Exceptions

NON-CROP SPRAYING

Canada thistle, White top and Meadow buttercup, (in non-crop areas such as roadsides, fence rows, rights-of-way and similar places): For spot treatment, use 3.0 ounces to 3.0 to 4.0 gallons of water or 0.5 gallon per acre in 12.0 to 20.0 gallons of water to give coverage for most extensive areas. Spray to wet weeds thoroughly when in bud to early bloom and again on fall regrowth.

Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

RESTRICTIONS AND LIMITATIONS FOR NON-CROP SPRAYING: For Broadcast Application: Do not apply more than 2.25 pints of product per acre per year (1.5 pounds acid equivalent per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

For spot treatment: Do not apply more than 4.6 pints of product per acre per year (3.0 pounds acid equivalent per acre per year).

Method

[Spot treatment](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

When in bud to early bloom and again on Fall regrowth.