

PEANUTS - ANNUAL MORNINGGLORY, ETC.

General Information

PRODUCT INFORMATION

Apply this product only as specified on the label.

NOTE: Avoid spray drift to susceptible plants and crops: cotton, tobacco, tomatoes, ornamentals, etc. Coarse sprays are less likely to drift.

Spray equipment previously used to apply another phenoxy must be thoroughly cleaned with alkali and water prior to use with 2,4-DB 175. Similarly, after using 2,4-DB 175, wash sprayer thoroughly before using again to spray susceptible crops.

Observe all caution and limitations on labeling of all products used in mixtures.

Use agitation to keep solution well mixed, especially if it has been allowed to stand unagitated.

Calibrate sprayer accurately.

Do not apply when crop is stressed from lack of moisture.

Do not apply this product through any type of irrigation system.

Do not use in or near greenhouses.

Limitations, Restrictions, and Exceptions

PEANUTS

Application:

Apply this product postemergent as an overall spray by ground sprayer or airplane. With ground applications, use a boom sprayer with flat fan-type nozzle. Adjust the height of the boom above the crop so the growing terminals of weeds are sprayed. Use low pressure (about 20 to 40 pounds per square inch). Do not use boom jet nozzles. With airplane applications, this product has been successfully applied using 5 gallons of total solution per acre. The height of the application should insure

thorough coverage of the tops of the weeds. Best control has resulted from airplane application with coarse spray when cocklebur is at least crop high and actively growing. Other weed species should be in the seedling stage and actively growing. Airplane application is not recommended in the vicinity of susceptible crops.

Do not apply if peanuts are suffering from lack of moisture.

Do not apply more than two (2) applications per season; second application should be made no later than the late bloom stage of peanuts (about 90 to 100 days after planting). Do not feed treated peanut vines or peanut hay to meat or dairy livestock. Do not apply to peanuts within 60 days of harvest.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

60 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

48 hours

Timings

[Postemergence \(Crop\)](#)

[Postemergence \(Weed\)](#)