

PREVENTION OF BROWN PATCH

General Information

DIRECTIONS FOR USE

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on turfgrass areas consisting of Kentucky bluegrass, annual bluegrass, bentgrass, perennial ryegrass, fine fescue, bermudagrass or mixtures where these grasses predominate.

Do not contaminate feed or food stuffs. Do not graze treated areas. Do not feed clippings to livestock. Keep children and pets off treated area until material has been washed off the foliage and the area is completely dry.

Precautions

Do not enter or allow others to enter the treated area (except those involved in watering-in) until watering-in is complete and the surface is dry.

Application Restrictions for Turfgrass Uses

Use on turfgrass is limited to golf course turf only. Use on turfgrass is prohibited on all other residential and commercial sites.

Minimum retreatment interval is 14 days.

Application to golf courses, including tees, greens, fairways, and roughs, is permitted only if the turfgrass is 2.5 inches or less in height.

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Refer the Suggested Spreader Settings.

Limitations, Restrictions, and Exceptions

PREVENTION OF BROWN PATCH

- Apply to moist or dry foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

•

Timings

[When environmental conditions favor disease activity.](#)