

BROCCOLI, ETC.

General Information

For the control of cutworms, armyworms, field crickets, house crickets, mole crickets, darkling beetles, grasshoppers, sowbugs and ants (except fire ants, harvester ants, pharaoh ants and carpenter ants).

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Carbaryl residues when Carbaryl is used in accordance with label directions.

If Carbaryl insecticide is used in accordance with label directions, carrot tops, sugar beet tops and corn forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

APPLICATION METHOD

This product may be applied only by ground using a mechanical spreader or underground shank applicator. Do not apply by hand, spoon or belly grinder. EXCEPTION: Aerial applications only to rangeland are allowed only under the auspices of the Animal and Plant Health Inspection Service (APHIS) Rangeland Grasshopper and Mormon Cricket Suppression Program, and/or allied state programs.

Limitations, Restrictions, and Exceptions

USE RATE

Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kohlrabi

Apply 40 lbs. per acre per application with a maximum yearly application of 120 lbs. per acre.

- Make applications only within 30 days of crop emergence or transplanting.

- Retreatment interval = 7 days.
- Do not apply within 3 days of harvest

Method

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

3 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[Within 30 days of crop emergence or transplanting.](#)