

FRUIT CROPS - CITRUS (SIX-SPOTTED MITES AND CITRUS RUST MITE)

General Information

USE PRECAUTIONS

IMPORTANT: Do not apply this product if temperatures during or within 3 days after application are expected to exceed 90°F in the area to be treated. Do not make applications within 2 weeks of an oil spray. Sulfur may burn foliage and fruit during periods of high temperature and under certain climatic conditions.

CALIFORNIA DRIFT RESTRICTIONS

The following conditions must be followed when using this product in the State of California for both ground and air applications.

1. The operator of the property and the applicator must establish a buffer zone of enough distance to prevent drift onto non-target areas such as hospitals, clinics, schools, residential areas and any other area designated by the county agricultural commissioner.
2. The operator of the property and the applicator must evaluate the method and equipment for each site to ensure proper and safe use. Evaluations shall include, but not be limited to the appropriateness of ground or air application.
3. Sulfur shall not be applied when the wind velocity exceeds 10 miles per hour. Applicators should be aware that in some areas, "dead calm" conditions are often associated with an inversion situation. In these areas, applying sulfur when there is a minimum air movement of 2 miles per hour will ensure that an inversion situation does not exist.
4. When applying sulfur dust, best application practices should be utilized. Contact your local pest control advisor or county cooperative extension advisor to obtain information regarding best application practices.

PRODUCT INFORMATION

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing and number of applications needed will vary with local conditions.

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground power equipment or aircraft. Except as otherwise shown, apply at the first sign of infection and repeat at 7-10 day intervals as needed to maintain control.

Limitations, Restrictions, and Exceptions

CITRUS

Avoid applications during or preceding high temperatures. Do not apply within 21 days of an oil spray.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Timings

[At the first sign of infection and repeat at 7-10 day intervals as needed to maintain control.](#)