

# **CURATIVE TREATMENT FOR BLACK LAYER**

## General Information

ECLIPSE may be applied at any time of the year as a preventative or curative treatment for Black Layer and to improve seed establishment when overseeding.

## Limitations, Restrictions, and Exceptions

## DIRECTIONS FOR USE

To achieve best results, and expose ECLIPSE to the Black Layer, mechanical aeration (hollow tine, solid tine, slitting, etc.) is crucial.

Aerate or spike to a depth greater than the Black Layer. Apply ECLIPSE at the recommended rate. Using drag brushes or other appropriate method to drag as much of the product into the aeration holes as possible. Leave aeration holes open, or if necessary, topdress using appropriate sized material. Until recovered from Black Layer, do not over irrigate, and refrain from using Iron, Sulfur or organic based products. Use nitrate based fertilizers and micronutrients like Manni-Plex products.

## APPLICATION RATES

Apply using conventional rotary or drop spreaders.

## BLACK LAYER

Curative Treatment: Broadcast at the rate of 6 lbs. per 1000 sq. ft., or approximately 5 bags per acre, following recommended pre-application procedures as shown above. Irrigate lightly after application. Up to 4 annual treatments may be made.

## Method

[Broadcast](#)

## Rates

[field rates 0](#)

[field rates 1](#)

-

Timings

Any time of the year.