

ROW CROPS

General Information

DIRECTIONS FOR USE

Use this product on soils to control various plant nematodes such as: root knot, root-lesion, citrus, stubby-root, soybean cyst, sting, lance, stylet and dagger.

NEO-TEC G.S.O. may be applied using any hand or power operated gravity or rotary type spreader designed for application of granular types of materials. NEO-TEC G.S.O. may be used for pre-planting of annual plants or post-planting applications of annual, perennial or established plants, trees, or turf areas. NEO-TEC G.S.O. is also suitable for treatment of individual trees and shrubs.

REPEAT APPLICATIONS

NEO-TEC G.S.O. should be applied at least twice during the spring and autumn seasons. If conditions warrant, NEO-TEC G.S.O. may also be reapplied on an as needed basis.

Neo-Tec G.S.O. is activated by the presence of water, either by rainfall or by artificial application.

Limitations, Restrictions, and Exceptions

ROW CROPS

Cotton, soybean, potatoes, sugar beets.

Band Dosage Rate: 5 lbs. per 1,000 ft. of row. This provides 67 lbs/planted acre based on 3 ft. rows, and 200 lbs/acre of planted band.

At Planting: Apply NEO-TEC G.S.O. in an 8-12 inch band along seed furrow. Seed depth dirt coverage is adequate for NEO-TEC G.S.O. Plant seed in or slightly below treated zone. POST-EMERGENCE: Side-dress NEO-TEC G.S.O. 4-6 inches each side of plant row and incorporate into the top 3 inches of soil. Repeat application as needed.

Method

[Band application](#)

Rates

[field_rates 0](#)

-

Timings

[At-Plant](#)

[Postemergence \(Crop\)](#)