

## **VEGETABLES**

### General Information

#### DIRECTIONS FOR USE

Use this product on soils to control various plant nematodes such as: root knot, root-lesion, citrus, stubby-root, soybean cyst, sting, lance, stylet and dagger.

NEO-TEC G.S.O. may be applied using any hand or power operated gravity or rotary type spreader designed for application of granular types of materials. NEO-TEC G.S.O. may be used for pre-planting of annual plants or post-planting applications of annual, perennial or established plants, trees, or turf areas. NEO-TEC G.S.O. is also suitable for treatment of individual trees and shrubs.

#### REPEAT APPLICATIONS

NEO-TEC G.S.O. should be applied at least twice during the spring and autumn seasons. If conditions warrant, NEO-TEC G.S.O. may also be reapplied on an as needed basis.

Neo-Tec G.S.O. is activated by the presence of water, either by rainfall or by artificial application.

#### Limitations, Restrictions, and Exceptions

#### VEGETABLES

Tomatoes, bell peppers, squash, strawberries, eggplants, cucumbers, carrot, radish, turnips, onions, peas, melons.

Broadcast Dosage Rate: 200 lbs. per acre. In spring or fall just prior to or during post harvest tilling, a broadcast application over entire field may be used at 200 lbs. of NEO-TEC G.S.O. per acre. NEO-TEC G.S.O. is to be incorporated into soil at one to four inches deep or lightly covered with soil. Repeat application as needed.

#### Method

[Soil incorporation](#)

[Broadcast](#)

Rates

[field\\_rates 0](#)

- 

Timings

[In spring or fall just prior to or during post harvest tilling.](#)