

POTATO

General Information

GENERAL INSTRUCTIONS

GROUND AND AERIAL APPLICATION: This product can be applied as a ground or aerial spray to control fungal infestations on listed crops. Application rates are for general use and must not be exceeded. The state agricultural extension or agricultural experiment station specialist should be consulted for specific applications and timing recommendations. With any spray application, thorough coverage is essential for good control. Do not apply this product through any type of irrigation system except on potatoes. Apply this product only as specified on this label. In case of accidental exposure, see First Aid Statement.

Do not allow this product to drift from the target site.

CHEMIGATION: Do not apply this product through any type of irrigation system on crops other than potatoes. Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. For specific information about calibration, contact State Extension Service specialists, equipment manufacturers or other irrigation experts.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustment should the need arise.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any

public areas such as schools, parks, playgrounds or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and symbol shall be a color, which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER. This sign is in addition to any sign posted to comply with the Worker Protection Standard.

SPRINKLER CHEMIGATION: The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a

positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

PRECAUTIONS: Nufarm does not recommend mixing with surfactants, spreaders, stickers or buffers unless testing or prior experience has shown the mixture to be non-phytotoxic to the crop. Combinations with some pesticides, micronutrients, spreaders, stickers, surfactants or buffering agents can increase phytotoxicity. Phytotoxicity may be severe. Emulsifiable concentrate insecticides can be especially injurious in combination. Do not graze dairy or meat animals in treated areas.

Limitations, Restrictions, and Exceptions

POTATO, Early Blight, Late Blight, Suppression of Colorado Potato Beetle. Apply 2.5 to 3.75* dry ounces per acre (15-10 acres/2.34 lb. per twin-pack or 7.5-5 acres per single pack). Ground (Closed Cabs Only): Apply in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. A spray pressure of less than 200 psi is recommended. Aerial: Apply in 3 to 10 gallons of water. For helicopter application, fly high enough so as not to whip the vines. Chemigation: Add the product through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended. Diluted spray should be directed uniformly to all parts of the plant and the gallonage increased according to the size of the plants. Application should begin with the appearance of blight weather conditions and continue on a 7-day schedule. Do not treat within 7 days of harvest. Do not exceed 11.25 ounces/acre per season. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area.

*PEST MANAGEMENT SYSTEMS. When used in combination with another fungicide registered for disease control on potatoes, a 1.87 ounce/acre rate may be employed.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[Application should begin with the appearance of blight weather conditions and continue on a 7-day schedule.](#)