

WOODY BRUSH AND TREES - SAGE, BLACK/SAGEBRUSH, CALIFORNIA/CHAMISE/TALLOWTREE, CHINESE

General Information

GENERAL INFORMATION

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THE LABEL OR CURRENT SUPPLEMENTAL LABELING ISSUED BY MANUFACTURER.

This product, a water soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

Hand-held sprayers may also be used.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of aboveground growth and deterioration of underground plant parts.

Unless otherwise specified on the label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of the label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Reduced or unacceptable control may result if weeds or brush are treated under poor growing conditions such as drought stress, disease or insect damage. Reduced results may also occur when treating weeds or brush heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, it is recommended that a residual herbicide program specified on the label be used. Read and carefully observe the precautionary statements and all other information appearing on the labels of all herbicides used.

Mixing this product with herbicides or other materials not recommended in the label may result in reduced performance. However, unless otherwise prohibited on the label or the label of an intended tank mix product, this product may be applied in combination with any herbicide registered for the same site, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. LIABILITY FOR CROP INJURY, HERBICIDE NONPERFORMANCE OR OTHER LOSS OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THE LABEL, OR SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT, IS SPECIFICALLY DISCLAIMED BY MANUFACTURER. BUYER AND ALL USERS ARE RESPONSIBLE FOR ALL LOSS OR DAMAGE IN CONNECTION WITH THE USE OR HANDLING OF MIXTURES OF THIS PRODUCT OR OTHER MATERIALS THAT ARE NOT EXPRESSLY RECOMMENDED IN THE LABEL.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs

and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

Limitations, Restrictions, and Exceptions

NOTE: If brush has been mowed or tilled or trees have been cut, treatment will not be effective until regrowth has reached the recommended stages of growth.

Apply this product when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See "DIRECTIONS FOR USE" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of the label for labeled uses and specific application instructions.

Apply this product as follows to control or partially control the following woody brush and trees.

SAGE, BLACK/SAGEBRUSH, CALIFORNIA/CHAMISE/TALLOWTREE, CHINESE – For control of these species, apply a 1 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

Tan oak resprouts – For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with fall applications.

Method

[Foliar spray](#)

[Hand-Held Spray](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[Postemergence \(Weed\)](#)