

ORNAMENTALS

General Information

GENERAL INFORMATION

This product is NOT recommended for application by homeowners.

In the state of New York, this product may be applied by commercial applicators ONLY.

This product controls crabgrass, other annual grasses and broadleaf weeds in established lawns, ornamental turf (including golf course fairways, roughs and tee boxes), and landscape ornamentals.

This product will not control emerged weeds, except for control of crabgrass up to initiation of tillering because the tillering stage of crabgrass growth generally corresponds to the time when seedlings are first visible in established turfgrasses.

Applying to crabgrass after tillering starts will not provide satisfactory control, and all other applications of this product should be made prior to germination of target weeds.

To be effective, this product must be activated by at least 1/2 inch of rainfall or irrigation. Applications should be timed to ensure that activation occurs prior to the tillering stage of crabgrass development or prior to germination of all other weeds.

USE PRECAUTIONS

- Do NOT allow the pouches to become wet prior to adding them to the spray tank.
- Do NOT remove the water-soluble pouches from the protective over-wrap except to add them directly to the spray tank.
- Do NOT handle the pouches with wet hands or wet gloves.
- Always reseal over-wrap bag to protect remaining unused pouches.

Limitations, Restrictions, and Exceptions

Note Regarding Application to Turf or Ornamental Species Not Listed on the label:

The user assumes the responsibility for any plant damage or other liability resulting from use of this product on species not recommended in the label.

Use Precautions

- Apply this product only to established ornamentals (after transplanting when soil around roots has been thoroughly settled by rainfall or irrigation).
- Do NOT apply this product on grasses grown for seed.
- Do NOT graze livestock or feed forage cut from areas treated with this product.
- Do NOT apply this product in enclosed structures and greenhouses.
- Do NOT apply this product through any type of irrigation system.
- Do NOT apply this product to the bare roots of ornamental plants.
- Do NOT incorporate this product into soil or dilution of the active ingredient and possible injury to plant roots may occur.
- Do NOT apply around ornamental plants that have been weakened or are under stress due to flooding, drought, wind injury, hail, frost damage, winter injury, excessive fertilizer or soil salts, injury due to insects, nematodes or diseases or injury from previously applied pesticides.
- Because drift or direct sprays of this product can injure the foliage of nontarget plants, do NOT make applications when weather conditions favor drift to non-target areas.

Maximum Use Rates

- Do NOT apply more than 0.5 lb. of product AI per acre (i.e., 4 water-soluble pouches per acre or 0.46 oz. per 1000 sq. ft.) per application, or more than 1.5 lbs. AI per acre per year (i.e., 12 water-soluble pouches per acre or 1.375 oz. per 1000 sq. ft.).

- In New York State, do NOT apply more than 0.25 lb. product AI per year (i.e., 2 water soluble pouches per acre or 0.23 oz. per 1000 sq. ft.).

Application Timing and Recommendations

Because the best weed control is obtained when applications are made to soil that is free of clods, weeds and debris such as leaves, prior to applying existing vegetation should be controlled by cultivation, hand weeding or with a postemergence herbicide.

Once this product is applied, do not disturb the soil surface as the herbicide barrier will be broken.

Care must be taken that soil or planting mixes have settled firmly following transplanting and that there are not cracks in the soil that would allow direct transmission of this product and plant roots.

Apple, Blueberry, Red maple, Maple Sugar, Mock Orange, Guava Pineapple, Japanese Quince, Rose, and Rosemary are Ornamental species only. DO NOT USE ON FOOD-PRODUCING TREES AND PLANTS.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

•

Restricted Entry Interval

12 hours

Timings

[Preemergence \(Crop\)](#)

[Preemergence \(Weed\)](#)