

MORE SUSCEPTIBLE WEEDS

General Information

USE RESTRICTIONS

Do not enter or allow entry until sprays have dried.

Do not apply this product through any type of irrigation system.

Do not use this product on or near desirable plants, including within the dripline of roots of desirable trees and shrubs, since injury may result.

Do not permit spray mist containing this product to drift onto them since even very small quantities of spray, which may not be visible, can cause severe injury during both growing and dormant periods.

Do not use the spray equipment for applying other materials to susceptible crops as injury may result.

USE PRECAUTIONS

Use care to avoid spray contact or drift to susceptible plants such as beans and other legumes, cotton, flowers, grapes, ornamentals, vegetables and other plants.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, air blast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Use only Medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a)

conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Additional Requirements for groundboom application:

Do not apply with a nozzle height greater than four feet above the target site.

WEEDS CONTROLLED

This product is safe to use on Washington, Toronto and Seaside bentgrasses, Kentucky and Merion bluegrasses. Apply this product on a warm, calm day when weeds are growing vigorously. Use any suitable sprayer or watering can with fine nose to give a uniform cover of the weeds. Although this herbicide is relatively slow in action, it gives a superior result on more difficult to control weeds.

TIME OF APPLICATION: To obtain the best results, this product must be applied when the weeds are most susceptible and conditions are such that no damage to the grasses will occur. Apply this product when the weeds are growing vigorously. During periods of drought, weeds become more resistant to herbicides. Under the same conditions, fine grasses are more susceptible to this product, particularly when temperatures exceed 90°F. This chemical is absorbed by the foliage of broadleaf weeds and is translocated throughout the whole plant to roots, etc. where it eventually causes death of the plant. Do not cut or mow the lawn 2 or 3 days before or after treatment so that maximum uptake and translocation can occur. Do not spray if rain is likely, otherwise the chemical may be washed off before absorption has taken place. For the same reason, do not water the lawn for at least 24 hours after application. Better control of the weeds classified as Less Susceptible Weeds may be obtained with a repeat application of half the recommended dose not more than 30 day interval.

This product is not recommended for St. Augustine grass and Centipedegrass, as it can cause damage to these species. Do not use this product on newly sown lawns or turf until grasses are well established. Reseed no sooner than three to four weeks after application of this product. Avoid drift onto other plants and do not use the first cuttings after treatment as a mulch for flowers and vegetables.

Limitations and Restrictions:

- For broadcast treatments, do not apply more than 3 pints of product per acre per application. Do not make more than 2 applications per year with a minimum of 30

days between applications.

- For spot treatments, do not apply more than 5 pints of product per acre per application. Do not make more than 2 applications per year with a minimum of 30 days between applications. Broadcast application is prohibited at this rate. A spot treatment is defined as a treatment area not greater than 1,000 square feet per acre.

Limitations, Restrictions, and Exceptions

TIME OF APPLICATION: To obtain the best results, this product must be applied when the weeds are most susceptible and conditions are such that no damage to the grasses will occur. Apply this product when the weeds are growing vigorously. During periods of drought, weeds become more resistant to herbicides. Under the same conditions, fine grasses are more susceptible to this product, particularly when temperatures exceed 90oF. This chemical is absorbed by the foliage of broadleaf weeds and is translocated throughout the whole plant to roots, etc. where it eventually causes death of the plant. Do not cut or mow the lawn 2 to 3 days before or after treatment so that maximum uptake and translocation can occur. Do not spray if rain is likely, otherwise the chemical may be washed off before absorption has taken place. For the same reason, do not water the lawn for at least 24 hours after application.

Better control of the weeds classified as Less Susceptible Weeds may be obtained with repeat applications of half the recommended dose carried out at not less than monthly intervals.

This product is not recommended for St. Augustine grass and Centipedegrass, as it can cause damage to these species. Do not use this product on newly sown lawns or turf until grasses are well established. Reseed no sooner than three to four weeks after application of this product. Avoid drift onto other plants and do not use the first cuttings after treatment as a mulch for flowers and vegetables.

Method

[Sprayer](#)

Rates

[field rates 0](#)

•

Timings

When weeds are growing vigorously.