

CANEBERRY

General Information

APPLICATION PROCEDURES

Apply by air or ground in sufficient water for complete coverage of the foliage. Use a minimum of 5 gallons of water for aerial application and 20 gallons by ground.

CROPS

NOTE: Mixing this product with certain surfactants, foliar fertilizers, or other pesticides can cause crop injury. Determine crop sensitivity to a particular combination by spraying a small area of foliage and fruit. Evaluate 3 to 7 days later for adverse effects.

NOTE: To assure the compatibility of this product with these and other products, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for 5 minutes. If the combination remains mixed, or can be re-mixed readily, the mixture is compatible.

NOTE: Mixing of this product with other products has been known to increase the salt content and the potential for fruit burn. Environmental factors that could exasperate burn potential include applying product during the following conditions: 1) temperatures above 90°F, 2) shortly after a rain event, 3) during color break of the fruit. Apply with extreme caution when these conditions exist. Determine crop sensitivity to these factors by spraying small areas of foliage and fruit. Evaluate 7 to 10 days later for adverse effects.

Limitations, Restrictions, and Exceptions

CANEBERRY

NOT FOR THIS USE IN CALIFORNIA.

Apply 4.5 pints product in a minimum of 20 gallons of water to assure thorough wetting of the foliage.

East of the Rocky Mountains: Begin application in the spring after bud break (1 to 3

inches of new growth) and repeat on 45 to 60-day intervals. Do not make more than 4 sprays during the growing season.

West of the Rocky Mountains: Fall Application – apply when conditions favor disease development and repeat if necessary in 3 to 4 weeks. Spring Application – make first application after bud break and repeat 3 to 4 weeks later.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[In new plantings, begin application when the plants produce new growth of 1 to 3 inches.](#)

[In established plantings, begin application when conditions favor disease development.](#)