

POTATOES - IN-FURROW APPLICATION

General Information

APPLICATION PROCEDURES

Apply by air or ground in sufficient water for complete coverage of the foliage. Use a minimum of 5 gallons of water for aerial application and 20 gallons by ground.

CROPS

NOTE: Mixing this product with certain surfactants, foliar fertilizers, or other pesticides can cause crop injury. Determine crop sensitivity to a particular combination by spraying a small area of foliage and fruit. Evaluate 3 to 7 days later for adverse effects.

NOTE: To assure the compatibility of this product with these and other products, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for 5 minutes. If the combination remains mixed, or can be re-mixed readily, the mixture is compatible.

NOTE: Mixing of this product with other products has been known to increase the salt content and the potential for fruit burn. Environmental factors that could exasperate burn potential include applying product during the following conditions: 1) temperatures above 90°F, 2) shortly after a rain event, 3) during color break of the fruit. Apply with extreme caution when these conditions exist. Determine crop sensitivity to these factors by spraying small areas of foliage and fruit. Evaluate 7 to 10 days later for adverse effects.

Limitations, Restrictions, and Exceptions

POTATOES

IN-FURROW APPLICATION: For the suppression of storage rot diseases such as Pink Rot (caused by *Phytophthora erythroseptica*) and Pythium Leak (caused by *Pythium* spp.) apply 3.75 to 10 pints product in combination with 11.6 to 13.0 fluid ounces of Ultra Flourish brand mefenoxam fungicide. Apply in a band at planting directly over the seed pieces prior to row closure in a minimum of 3 gallons of water per acre.

Additional in-season foliar applications of this product tank mixed with an appropriate fungicide containing either mefenoxam, chlorothalonil, mancozeb, or copper is necessary when:

- disease conditions are severe or
- the variety planted is susceptible or moderately susceptible to the disease or
- field is located in a long growing season location

See Ultra Flourish Specimen Label and Ultra Flourish Supplemental Label for more information.

Method

[In Furrow](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[At-Plant](#)