

TOMATO

General Information

APPLICATION PROCEDURES

Apply by air or ground in sufficient water for complete coverage of the foliage. Use a minimum of 5 gallons of water for aerial application and 20 gallons by ground.

CROPS

NOTE: Mixing this product with certain surfactants, foliar fertilizers, or other pesticides can cause crop injury. Determine crop sensitivity to a particular combination by spraying a small area of foliage and fruit. Evaluate 3 to 7 days later for adverse effects.

NOTE: To assure the compatibility of this product with these and other products, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for 5 minutes. If the combination remains mixed, or can be re-mixed readily, the mixture is compatible.

NOTE: Mixing of this product with other products has been known to increase the salt content and the potential for fruit burn. Environmental factors that could exasperate burn potential include applying product during the following conditions: 1) temperatures above 90°F, 2) shortly after a rain event, 3) during color break of the fruit. Apply with extreme caution when these conditions exist. Determine crop sensitivity to these factors by spraying small areas of foliage and fruit. Evaluate 7 to 10 days later for adverse effects.

Limitations, Restrictions, and Exceptions

TOMATO

Apply 2.5 to 5 pints product in a minimum of 10 gallons of water per acre beginning at the 2 to 4-leaf growth stages for direct seeded tomatoes or immediately after transplanting to the field. Continue on a 7 to 14-day spray interval when conditions are favorable for disease development. Use the higher rate (3.75 to 5 pints product in a minimum of 10 gallons of water per acre) and shorter spray interval when disease pressure is moderate to high.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[Beginning at the 2 to 4-leaf growth stages for direct seeded tomatoes or immediately after transplanting to the field.](#)