

# **SORGHUM: FOR END-USE APPLICATION AT AGRICULTURAL ESTABLISHMENTS**

## General Information

### DIRECTIONS FOR USE

For use in commercial seed treaters only, with the exception of application to canola, cotton (delinted seed), field corn, sorghum, wheat and barley, which may be made either by commercial seed treatment or as an end-use seed treatment on agricultural establishments at, or immediately before, planting. This product is to be used in liquid or slurry treaters.

### USE RESTRICTIONS

Rape greens grown and harvested from this product's treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from this product's treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.

Wheat, Barley, Oats, Rye, Triticale, Sorghum: Do not graze or feed livestock on treated areas for 45 days after planting.

Cotton (delinted seed only): Regardless of the type of application (seed treatment, soil or foliar) do not apply more than a total of 0.5 pound of imidacloprid per acre per cropping cycle.

Stored Seed Protection: Applied at labeled rates equal to or above 0.8 fluid ounce per hundredweight, this product will provide protection to seed against injury from the following insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryzae*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). It is recommended that seed with existing populations of stored grain pests be fumigated prior to treating and bagging seed.

For Early Season Protection Against Certain Sucking Insects: This product will aid in the protection of seeds and seedlings against injury by certain early season insects.

### Limitations, Restrictions, and Exceptions

## SORGHUM

For End-Use Application at Agricultural Establishments: Apply using an HCBT or an Eight-Bag Batch Treater. Shake this product thoroughly before use. Dilute 3.2 fluid ounces of this product with water. The final slurry rate should be adjusted to apply a rate of 8 to 10 fluid ounces of diluted slurry per 50-pound bag of seed. Treat one-half of seed with one-half of slurry mix. Add the balance of the seed and apply balance of slurry. Allow mixing until seed is thoroughly covered. Apply 0.75 ounce of dry TALC per 50-pound bag of seed following the product application and allow it to distribute evenly on the seed. Do not graze or feed livestock on treated areas for 45 days after planting.

### Method

[Seed Treatment](#)

### Rates

[field\\_rates 0](#)

- 

### Restricted Entry Interval

12 hours

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

### Timings

[N.A.](#)