

COMMERCIAL COTTON SEED TREATMENT

General Information

DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Limitations, Restrictions, and Exceptions

COMMERCIAL COTTON SEED TREATMENT

This product is recommended for use at time of planting as a seed treatment to protect against sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*) which impairs good cotton seed germination and seedling development. This product can be tank-mixed with other properly labeled cottonseed protectants for *Pythium* control. This product is a liquid formulation that can be pumped or poured directly into the cotton seed treater or diluted with water, provided the recommended amount of product is applied.

Use 1.25 to 4.0 fluid ounces of this product per 100 pounds cottonseed to reduce seedling damage caused by sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*).

Label Treated Seed: "This seed is treated with Spera™ 240 FS Fungicide, which contains the active ingredient myclobutanil. Do not use for food, feed, oil production or any other purpose except planting. Do not graze cotton plants grown from treated seed. Store away from feed and foodstuffs."

Always use this product with an EPA approved dye or colorant that imparts an unnatural color to the seed.

Method

[Seed Treatment](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At the time of planting.](#)