

GOLF COURSES: TEES, GREENS AND ORNAMENTAL TURF - (PREVENTIVE - DOLLAR SPOT, LARGE BROWN PATCH)

General Information

GENERAL INFORMATION:

TM + CTN E-Pro 66.6 WDG Fungicide is a formulation of two chemicals that provides broad spectrum contact and systemic disease control on golf courses and ornamental turfgrass as well as on ornamental plants, shrubs and trees when used according to the directions on the label.

Use Precautions

- Use of this product on home lawns is prohibited.
- Do NOT graze treated areas.
- Do NOT feed clippings to livestock or poultry.

General Mixing and Application Instructions

1. Add the recommended amount of TM + CTN E-Pro 66.6 WDG Fungicide to the spray tank while filling with water.
2. Maintain continuous agitation when filling the spray tank and during spray operations.
3. For best disease control results, thorough and uniform coverage of the plant surfaces is essential.

NOTE: Tank mixing this product with surfactants, fertilizers, and / or other pesticides is not recommended unless prior use has shown the combination to be physically compatible, effective and non-injurious under your conditions of use.

- Do NOT apply this product through any type of irrigation system.

- Do NOT apply within 150 ft. (for air-blast) or 25 ft (for ground applications) of marine / estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the body of water.

Limitations, Restrictions, and Exceptions

GOLF COURSES: TEES, GREENS AND ORNAMENTAL TURF - (PREVENTIVE - DOLLAR SPOT, LARGE BROWN PATCH):

Application Interval: 14 - 21 days

TM + CTN E-Pro 66.6 WDG Fungicide should always be used in conjunction with good turf management practices and programs.

Use Precautions

- After application, do NOT mow or water until spray deposit on turfgrass is thoroughly dry.
- Do NOT graze animals on treated turf.
- Do NOT feed clippings to livestock or poultry.
- Do NOT apply with fixed wing or rotary aircraft.

MAXIMUM APPLICATION RATES FOR THE PRODUCT

Greens

Maximum Seasonal Application Rate: 130.9 lbs / acre (48.08 oz / 1000 sq ft)

Maximum Individual Application Rate: 22.6 lbs /acre (8.3 oz / 1000 sq ft)

Tees

Maximum Seasonal Application Rate: 104 lbs / acre (38.2 oz / 1000 sq ft)

Maximum Individual Application Rate: 22.6 lbs /acre (8.3 oz / 1000 sq ft)

Fairways (Except Florida)

Maximum Seasonal Application Rate: 32.7 lbs / acre (12.0 oz / 1000 sq ft)

Maximum Individual Application Rate: 22.6 lbs /acre (8.3 oz / 1000 sq ft)

Fairways (Florida) (Make only 1 application at the maximum application rate)

Maximum Seasonal Application Rate: 16.3 lbs / acre (6.0 oz / 1000 sq ft)

Maximum Individual Application Rate: 16.3 lbs /acre (6.0 oz / 1000 sq ft)

Public Turf Areas (Make 4 applications per season)

Maximum Seasonal Application Rate: 52 lbs / acre (19.1 oz / 1000 sq ft)

Maximum Individual Application Rate: 16.3 lbs /acre (6.0 oz / 1000 sq ft)

For severe disease pressure, use the highest recommended rate and the shortest application interval listed. Note: Do NOT apply at less than a 14-day interval.

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[prior to disease development](#)