

SUGARCANE - SECOND APPLICATION PER SEASON - JOHNSONGRASS

Limitations, Restrictions, and Exceptions

SUGARCANE - SECOND APPLICATION PER SEASON - JOHNSONGRASS

A second application may be made 4 to 10 weeks after the first application if needed to control weeds prior to closure of the planted sugarcane.

(Use in heavy weed infestations or re-infestation or where weed species germinate at different times in the same growing season)

RESTRICTIONS AND PRECAUTIONS: Do not rotate to any crop which is not registered for use with products containing asulam for one year following the last application of this product.

Cover crops may be planted if they are plowed under and not grazed; otherwise, do not plant crops other than sugarcane in treated areas.

Do not graze or feed sugarcane fodder and forage to livestock.

Asulam 3.3 Herbicide should be used when weeds are actively growing and may be applied to plant cane or to cane grown from stubble.

Do not harvest sugarcane within the following number of days after application: 140 days in mainland USA (excluding Louisiana), 100 days in Louisiana and 400 days in Hawaii.

Differences in crop tolerance to asulam among sugarcane varieties have been reported in Louisiana. Contact your local County Agent or University Extension Specialist for further information.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

-

Timings

[When the grass is 18 - 24 inches tall.](#)