

CORN - POSTEMERGENCE - BROADLEAF AND GRASS WEED CONTROL

General Information

GENERAL INFORMATION AND RESTRICTIONS

Atrazine 4L herbicide will control many annual broadleaf and grass weeds in corn, sorghum, sugarcane and certain other crops specified on the label.

WEEDS

Atrazine may be applied before or after weeds emerge. The continuous use of this product and other triazines may cause the development of biotypes of grasses and broadleaf weeds that cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes, Atrazine 4L should be either tank mixed or used in sequence with a herbicide approved for use with triazine which is not in the triazine class of chemicals. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide. Read and follow all directions for use and precautions on the labels of products to be tank mixed with atrazine. Consult your state Extension Service for specific recommendations for your area.

In each case where a range of rates is given, the lower rate should be used on coarse-textured soils and soils low in organic matter, and the higher rate should be used on fine-textured soils and soils high in organic matter. Since atrazine acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control. This product is nonflammable, noncorrosive to equipment and metal surfaces, and has low electrical conductivity.

Do not apply this product through any type of irrigation system.

When tank-mixing or sequentially applying atrazine or products containing atrazine to corn or sorghum, the total pounds of atrazine applied (lbs ai/A) must not exceed 2.5 lbs ai/A/year. When tank-mixing or sequentially applying atrazine or products

containing atrazine to corn or sorghum, do not exceed an application rate of 2.0 pounds active ingredient of atrazine per acre for any single application and the total pounds of atrazine applied (lbs ai/A) must not exceed 2.5 pounds active ingredient per year.

When tank-mixing or sequentially applying atrazine or products containing atrazine to crops other than corn or sorghum, the total pounds of atrazine applied (lbs ai/A) must not exceed the specific seasonal rate limits as noted in the use directions.

Care should be taken to avoid using atrazine where adjacent desirable trees, shrubs or plants may be injured. Avoid using near adjacent desirable plants or in greenhouses, or injury may occur.

Note: Winfield Solutions, L.L.C does not recommend the use of Atrazine 4L in combination with other herbicides or oils, except as specifically described on the label.

APPLICATION PROCEDURES

Ground Application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. For the most uniform distribution of broadcast applications, use 80 degrees flat fan-type nozzles. For band applications, use flat fan even-spray nozzles. Be certain that nozzles are uniformly spaced and are of the same size and type. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Screens and strainers should be no finer than 50 mesh between the pump and boom. Screens on the suction side of the pump should be 16 mesh or coarser. Do not place a screen in the recirculation line.

Use a pump with capacity to: (1) provide sufficient hydraulic agitation during mixing and application to keep the material in suspension, (2) maintain 30-40 psi operating pressure, and (3) provide a minimum of 20% by-pass at all times. (At minimum, 10 gallons/minute per 100 gallon tank size). Unless otherwise specified, use a minimum of 5-10 gallons of water per acre for preplant incorporated, pre-emergence and post (with or without oil or surfactant) applications.

Aerial Application: For preplant and pre-emergence broadcast treatments, apply at a 1:1 ratio Atrazine 4L to water (Example: Recommendation calls for 1 quart of product and is mixed with 1 quart of water) to be applied per acre. For

postemergence treatments on corn and sorghum, apply the recommended rate of Atrazine 4L plus oil in a minimum of 2 gallons of water per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. The above water volumes are the minimum to be applied per acre.

Atrazine 4L in Water Applications: Atrazine 4L, a liquified formulation, should be mixed with water and applied as a spray. Pour Atrazine 4L into the tank during or after filling. Hydraulic (jet) or mechanical agitation is recommended during mixing and application to keep the material in suspension. All return lines to the tank must discharge below liquid level and agitation should not be so violent as to cause air bubbles to form in the liquid. Wash sprayer thoroughly after use.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

CORN:

ATRAZINE 4L ALONE: Apply either before planting, at planting or after planting as indicated below. For soil applications prior to crop emergence (i.e., early preplant, preplant incorporated, preplant surface, at planting or preemergence) see Table 1.

Postemergence Broadleaf and Grass Weed Control with Atrazine 4L plus Emulsifiable Oil or Oil Concentrate in Water: Broadcast 4 pints per acre after weed emergence, but before weeds reach 1.5 inches in height and before corn reaches 12 inches in height. Add the following volume of one of the type oils indicated for aerial or ground application unless the oil label specifies otherwise.

Note: Crop-derived or petroleum-derived oil concentrates should contain at least 1%, but no more than 20% suitable emulsifier or surfactant blend. Petroleum-derived oil should contain at least 1% suitable emulsifier.

Precautions for Applications of Atrazine 4L plus Emulsifiable Oil in Water to Corn:

(1) Do not use oils in sprays when treating inbred lines or any breeding stock as severe injury may occur.

(2) Adding other insecticides, herbicides, liquid fertilizers or other materials is not recommended with atrazine and emulsifiable oil in water because they may cause

compatibility problems or crop injury.

(3) Store and handle emulsifiable oil carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.

(4) Do not use oil in atrazine sprays when crop is under stress from prolonged cold, wet weather, poor fertility, or other factors, or when crop is wet and succulent from recent rainfall, as crop injury may occur.

(5) Do not exceed 2.5 lbs. active ingredient (or 5 pts. of this product) per acre per calendar year.

(6) Postemergence application to corn must be made before corn exceeds 12 inches in height.

Precautions for all Applications to Corn:

(1) The maximum application rate for corn, including tank mixtures and sequential treatments, is 2.5 lbs. active ingredient per acre per calendar year. Therefore, do not apply more than 5 pts. of Atrazine 4L on corn in any one year.

(2) Following harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in the fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of the rate used.

(3) For best control of velvetleaf and cocklebur, the application rate cannot be less than 2 pounds per acre of active ingredient, either alone or in tank mix combinations.

(4) Do not graze treated areas for 21 days following application.

(5) Field Corn: Do not harvest for forage within 60 days of application.

(6) Sweet Corn: Do not harvest for forage within 45 days of application.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Postemergence \(Crop\)](#)

[Postemergence \(Weed\)](#)