

SORGHUM AND SORGHUM-SUDAN HYBRIDS (GRAINS AND FORAGE TYPES) - PREPLANT INCORPORATION, ETC.

General Information

GENERAL INFORMATION

This herbicide controls many annual broadleaf and grass weeds in corn, sorghum, sugarcane and certain other crops specified on the label. This product may be applied before or after weeds emerge. FOLLOWING MANY YEARS OF CONTINUOUS USE OF THIS PRODUCT AND CHEMICALLY RELATED PRODUCTS, BIOTYPES OF SOME OF THE WEEDS LISTED ON THE LABEL HAVE BEEN REPORTED WHICH CANNOT BE EFFECTIVELY CONTROLLED BY THIS AND RELATED HERBICIDES. WHERE THIS IS KNOWN OR SUSPECTED, WE RECOMMEND THE USE OF THIS PRODUCT IN COMBINATION WITH OTHER REGISTERED HERBICIDES WHICH ARE NOT TRIAZINES. CONSULT WITH YOUR STATE AGRICULTURAL EXTENSION SERVICE FOR SPECIFIC RECOMMENDATIONS.

When tank-mixing or sequentially applying atrazine or products containing atrazine to corn or sorghum, do not exceed an application rate of 2.0 pounds active ingredient of atrazine per acre for any single application and the total pounds of atrazine applied (lbs ai/A) must not exceed 2.5 pounds active ingredient per acre per year. When tank-mixing or sequentially applying atrazine or products containing atrazine to crops other than corn or sorghum, the total pounds of atrazine applied (lbs ai/A) must not exceed the specific seasonal rate limits as noted in the use directions.

APPLICATION PROCEDURES

Aerial application of dry flowable products that are not packaged in water soluble bags is prohibited. Ground Application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture per acre for all preplant, preemergence and postemergence applications (with or without oil or surfactant) with ground equipment.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

SORGHUM AND SORGHUM-SUDAN HYBRIDS (Grain and Forage Types)

Apply before planting, at planting, or after planting, as indicated in the directions below. See Corn section for weeds controlled.

Preplant Incorporated (Broadleaf and grass control): Broadcast in spring after plowing at rate in Table 3. Apply before, during, or after final seedbed preparation. Avoid deep incorporation. For best results, apply within two weeks before planting.

Preemergence (Broadleaf and grass control): Apply at rate shown in Table 3 during or shortly after planting and before weed or crop emergence.

Broadleaf and Grass Weed Control in Sorghum*

FOR ALL SOIL APPLICATIONS PRIOR TO CROP EMERGENCE

On Highly Erodible Soils

If conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues at planting, apply 2.2 lbs./A of Atrazine 90DF (2 lbs. ai/A maximum) as a broadcast spray.

If the soil coverage with plant residue is less than 30% at planting, a maximum of 1.8 lbs./A of Atrazine 90DF (1.6 lbs ai/A) may be applied.

On Soils Not Highly Erodible

Apply 2.2 lbs./A of Atrazine 90DF (2 lbs ai/A maximum) as a broadcast spray.

*Do not apply preplant in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN, or TX. Do not apply preemergence in NM, OK, or TX, except in northeast OK, and the TX Gulf Coast.

In case of planting failure, sorghum or corn may be replanted. Do not make second broadcast application, as injury may occur. If originally applied in a band and sorghum or corn is replanted in untreated row middles, this product may be applied in a band to the second planting. However, the maximum application rate for sorghum or corn is 2.8 lbs. per acre per calendar year.

Precautions for All Applications to Sorghum: (1) Heavy rain immediately following application, tends to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Do not apply to furrowplanted sorghum until furrows are leveled (plowed in). Level deep planter marks or seed furrows before application. (2) Application to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soil may result in crop injury. (3) Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used. (4) Injury may occur if both this herbicide, preplant or preemergence, and an atplanting systemic insecticide applied in-furrow are used. (5) Do not apply more than 2.5 lbs. active ingredient (2.8 lbs. product) per acre per year. (6) For all tank mixtures and sequential treatment of products containing atrazine, do not exceed 2.0 lbs. a.i./A of atrazine as a pre or post application or 2.5 lbs. a.i./A as the total of pre plus post applications per calendar year. (7) For all soil application prior to crop emergence (except for preemergence use on bedded sorghum in AZ and CA), do not apply on coarse textured soils; i.e., sand, loamy sand, sandy loam or to medium and fine textured soils having less than 1% organic matter, or crop injury may occur. (8) For postemergent applications, do not apply to sand or loamy sand, as crop injury may occur.

Note: Do not harvest for forage within 60 days of a pre-emergence application and 45 days of a post-emergent application.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[At-Plant](#)

[Preplant](#)

[Preplant Incorporated](#)

[Preemergence \(Weed\)](#)