

BLUEBERRIES, CANEBERRIES, CRANBERRIES AND STRAWBERRY

General Information

GENERAL INFORMATION

FOLI-GRO is manufactured specifically for use as a liquid foliar or soil applied fertilizer. Its use is suggested as part of a Total Nutrition System program to enhance yields and quality.

Under adverse environmental conditions (water stress, cold soils, salts, etc.) when used as directed FOLI-GRO may supply otherwise unavailable plant nutrients. Research has shown that foliarly applied nutrients are absorbed more efficiently by plants than are those supplied in the soil under the above conditions.

FOLI-GRO is uniquely formulated to supply nutrients in a readily available form for the plant to obtain a quick response to needed elements. A tissue testing program may be helpful to monitor and maintain optimum plant growth and quality. Nutrient translocation inside the plant is generally more rapid when applied foliarly.

DIRECTIONS FOR USE

FOLIAR APPLICATIONS (Ground or Air): Apply the recommended amount of FOLI-GRO Manganese 5% per acre per application throughout the growing season as needed to maintain adequate levels for growth. Severe deficiencies may require more frequent applications. Use sufficient water to insure good coverage. Do not exceed one quart per gallon of water. If applied too concentrated, slight burning of the foliage may occur.

Limitations, Restrictions, and Exceptions

BLUEBERRIES, CANEBERRIES, CRANBERRIES AND STRAWBERRY

Do not apply after fruit set.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)