

OTHER CROPS

General Information

GENERAL INFORMATION

MORA-LEAF Alfalfa Gold is formulated primarily for foliar applications to prevent or correct sulfur and boron deficiencies in a wide range of agronomic and ornamental plants. Its use is suggested as a supplement to a regular, balanced fertilizer program to enhance yields and improve quality.

Application of MORA-LEAF Alfalfa Gold is a means of obtaining a quick response to needed elements. Foliar applications of MORA-LEAF Alfalfa Gold may be particularly beneficial during periods of peak nutrient demand, for crops grown on soils having poor nutrient availability or to crops suffering from a weakened root system.

Avoid foliar nutrient applications to crops during periods of moisture, salinity or heat stress. Applications of foliar nutrients to sensitive crops such as leafy vegetables or soft-skinned fruits during periods of excessive temperatures may cause cosmetic injury.

MORA-LEAF Alfalfa Gold should be used as part of a comprehensive Total Nutrition System for optimizing plant growth, development, yield and quality.

DIRECTIONS FOR USE

MORA-LEAF Alfalfa Gold may be applied as a foliar spray by air or by ground. Maintain agitation to provide uniform distribution within the spray tank. Thorough and even spray coverage is essential for maximum nutrient uptake and crop safety. Always include a high quality wetting agent, such as R-11 or Sylgard 309, at recommended rates.

Optimum rate of application will vary based on a variety of factors including degree of need or deficiency, growth stage, yield potential, environmental conditions, existing soil fertility levels, etc.

Add MORA-LEAF Alfalfa Gold to spray tank that is at least 50% full with water and maintain agitation. MORA-LEAF Alfalfa Gold should dissolve rapidly and distribute uniformly. When water temperatures are cold, allow extra time for this product to

completely dissolve. Do not exceed 2 pounds MORA-LEAF Alfalfa Gold per gallon of water. Unless otherwise specified on tank mixed pesticide labels, add pesticide last. When combining with products packaged in water-soluble pouches, allow pouches to dissolve completely before adding nutrients.

Limitations, Restrictions, and Exceptions

OTHER CROPS

Use 5-10 pounds per acre in a program of multiple applications throughout the growing season, depending on the crop.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Timings

[N. A.](#)