

## **SUNFLOWERS (EXCEPT IN CALIFORNIA)**

### General Information

#### APPLICATION

Dilute DELTA GOLD 1.5 EC in water according to the directions and apply to the target crops at the rates listed. Uniform coverage is essential. Consult your local or state agricultural authorities regarding economic thresholds and timing of application for specific pests. Under heavy pest populations use of the highest labeled rate is recommended.

Do not apply Delta Gold 1.5EC by aerial application in New York State.

DELTA GOLD 1.5 EC may be applied before, during, or after planting. Use the higher labeled rate for soil incorporated applications.

#### RESISTANCE

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

#### Limitations, Restrictions, and Exceptions

#### SUNFLOWERS (EXCEPT IN CALIFORNIA)

##### Minimum Gallonage Requirements:

For aerial applications use a minimum of 2 gallons of water per acre. For ground

applications use a minimum of 5 gallons of water per acre for thorough coverage of the foliage.

NOTES for Use on Sunflowers:

- DO NOT graze or feed treated sunflower foliage to livestock.
- DO NOT apply this product to sunflowers as an Ultra Low Volume (ULV) spray.
- Cabbage looper, Corn earworm: For best control apply to 1st and 2nd instar larvae.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

[field\\_rates 2](#)

•

Restricted Entry Interval

12 hours

Timings

[N.A.](#)