

POME FRUITS - APPLE LEAFROLLER, APPLE LEAFMINER, ETC.

General Information

APPLICATION

Dilute DELTA GOLD 1.5 EC in water according to the directions and apply to the target crops at the rates listed. Uniform coverage is essential. Consult your local or state agricultural authorities regarding economic thresholds and timing of application for specific pests. Under heavy pest populations use of the highest labeled rate is recommended.

Do not apply Delta Gold 1.5EC by aerial application in New York State.

DELTA GOLD 1.5 EC may be applied before, during, or after planting. Use the higher labeled rate for soil incorporated applications.

RESISTANCE

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Limitations, Restrictions, and Exceptions

POME FRUITS

For applications to boring insects, apply prior to the insects entering the wood or fruit. Insects contained within plant tissue may not be controlled.

Minimum Gallonage Requirements:

For aerial applications use a minimum of 10 gallons of water per acre. For ground applications, use a minimum of 25 gallons of water per acre for thorough coverage of the foliage.

NOTES for Use on Pome Fruits:

- Allow 7 days between applications of DELTA GOLD 1.5EC
- Tarnished plant bug, Western plant bug: See resistance statement under the "Directions For Use" section.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

•

Restricted Entry Interval

12 hours

Timings

[Boring insects: Prior to the insects entering the wood or fruit.](#)