

BARLEY

General Information

SPRAY VOLUME

ONSET 3.6L may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Limitations, Restrictions, and Exceptions

BARLEY

Notes and Restrictions:

- Apply ONSET 3.6L in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.
- Straw cut after harvest may be fed or used for bedding.
- Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of ONSET 3.6L.
- Barley fields should be observed closely for early disease symptoms, particularly

when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.

General Comments:

- For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with ONSET 3.6L.
- ONSET 3.6L must have 2 to 4 hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, ONSET 3.6L will be resistant to weathering.
- ONSET 3.6L is a demethylation inhibitor (DMI) fungicide (Group 3).

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[Rusts: At the earliest sign of rust pustules on foliage.](#)

[Fusarium head blight: When main stem heads have fully emerged \(Feekes 10.5\) on 50% of the plants.](#)