

CORN - ANNUAL BROADLEAF WEEDS (POSTEMERGENCE)

General Information

GENERAL INFORMATION

PHENOXY 088 is a special formulation containing 2,4-D acid. It is more effective than 2,4-D amines for controlling hard-to-kill weeds such as field bindweed, Russian knapweed, Canada thistle, leafy spurge, cattails, tules and nutgrass. PHENOXY 088 should be applied when daytime temperatures are lower than 80 to 85 degrees. Spraying when temperatures are higher may cause rapid foliar burn and decrease the product's effectiveness. Treatment at higher temperatures also increases the possibility of crop injury. **INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.** Crop varieties vary in response to 2,4-D and some are easily injured. Apply PHENOXY 088 only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company or state Agricultural Extension Service for advice.

GENERAL RESTRICTIONS: Do not apply product through any type of irrigation system. Do not use in or near a greenhouse.

PHENOXY 088 contains 0.35 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.75 pounds of a.e. per acre per crop cycle.

Limitations, Restrictions, and Exceptions

- Annual Weeds - Beggarticks, common broomweed, knotweed, mallow (venice or little), musk thistle, wild carrot, vervains, smartweeds (annual species), salsify (western or common), russian thistle, prickly lettuce,: These species may require repeat applications and/or use of higher rate recommended on this product label even under ideal conditions for application.

- Pigweeds (*Amaranthus* spp.)- Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

- Mustards (except blue mustard)
- Pepperweeds (except perennial)

CORN RESTRICTIONS:

- Postemergence: Make no more than one application per crop cycle.
- Postemergence: Do not apply more than 1-1/3 pints per acre per application.
- Minimum spray interval between applications for sweet corn is 21 days.
- Do not use treated crop as fodder for 7 days following application.
- Corn (Field and Pop) Pre-Harvest Interval is 7 days.
- Corn (Sweet) Pre-Harvest Interval is 45 days.

(PHENOXY 088 contains 0.35 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 3.0 pounds of a.e. per acre per year for Field and Pop Corn.)

- Do not exceed a combined total of 1.5 pounds of a.e. per acre for Sweet Corn.
- A minimum application volume of 5 gallons per acre by air or 10 gallons per acre by ground is recommended.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[Postemergence \(Weed\)](#)

Apply when weeds are small and corn is less than 8 inches tall (to top of canopy).