

FRUIT AND NUT CROPS - OLIVES

General Information

GENERAL INFORMATION

Apply this herbicide before weeds emerge or after removal of weed growth. SIMAZINE controls a wide variety of annual broadleaf and grass weeds when used at selective rates in agricultural crops and ornamental plantings. When used at higher nonselective rates in noncrop areas, it also controls many perennial broadleaf and grass weeds.

Where a range of application rates is given, use the low rate on coarser textured soil and soil low in organic matter; use the high rate on fine textured soil and soil high in organic matter.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Since this product enters weeds mainly through their roots, moisture is needed to move it into the root zone. Very dry soil conditions and lack of rainfall following application may necessitate shallow cultivation.

This product is noncorrosive to equipment, nonflammable, and has low electrical conductivity.

Failure to follow the directions for use and precautions on the label may result in crop injury, poor weed control, and/or illegal residues.

AERIAL APPLICATION IS PROHIBITED.

FOLLOWING MANY YEARS OF CONTINUOUS USE OF SIMAZINE AND CHEMICALLY RELATED PRODUCTS, BIOTYPES OF SOME OF THE WEEDS LISTED ON THE LABEL HAVE BEEN REPORTED WHICH CANNOT BE EFFECTIVELY CONTROLLED BY THIS AND RELATED HERBICIDES. WHERE THIS IS KNOWN OR SUSPECTED AND WEEDS CONTROLLED BY THIS PRODUCT ARE EXPECTED TO BE PRESENT ALONG WITH RESISTANT BIOTYPES, WE RECOMMEND THE USE OF THIS PRODUCT IN REGISTERED COMBINATIONS OR IN SEQUENCE WITH OTHER REGISTERED HERBICIDES WHICH

ARE NOT TRIAZINES. IF ONLY RESISTANT BIOTYPES ARE EXPECTED TO BE PRESENT, USE A REGISTERED NON-TRIAZINE HERBICIDE. CONSULT YOUR STATE AGRICULTURAL EXTENSION SERVICE FOR SPECIFIC RECOMMENDATIONS.

Refer to the label for tank mix information.

Limitations, Restrictions, and Exceptions

FRUIT AND NUT CROPS

Apply the spray to the orchard or vineyard floor avoiding contact with fruit, foliage, or stems. Recommended rates are based on broadcast treatment. For band application, or spot applications around fruit or nut trees, reduce the broadcast rate of this product and water per acre in proportion to the area actually sprayed.

Use Precautions and Restrictions (All fruit and nut crops): To avoid crop injury: 1) Apply only to orchards or groves where trees have been established 12 months or more unless specified differently. 2) Make only one application per year, except as noted otherwise. 3) Do not use on gravelly, sand, or loamy sand soil. 4) Immediately following application, limit overhead sprinkler irrigation to ½ inch.

OLIVES

Apply 2.2-4.4 pounds per acre following grove preparation in the fall. Repeat annually in mid-winter. Do not apply more than 4.4 pounds per acre per calendar year. Do not apply more than once per calendar year.

Method

[Broadcast/Foliar Ground](#)

[Band](#)

[Broadcast/Foliar Ground](#)

[Band](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection

Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Preemergence \(Weed\)](#)

[In fall.](#)