

TURFGRASS - TO CONTROL SEVERE LEAF SPOT/MELTING OUT, ETC.

General Information

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not enter or allow others to enter the treated area until dusts have settled.

GENERAL INFORMATION

For use on turfgrass areas consisting of bentgrass, bluegrass, *Poa annua*, bermudagrass, St. Augustinegrass, ryegrass, fine fescue or tall fescue, or mixtures where these grasses predominate.

Controls these diseases while fertilizing turfgrasses: red thread, dollar spot, brown patch, pink snow mold (*Microdochium nivale*), leaf spot/melting out, anthracnose.

Suppression of these diseases: summer patch, necrotic ring spot, Fusarium blight

GENERAL PRECAUTIONS AND RESTRICTIONS

Not for homeowner use. For use by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products.

Use of this product at residential sites is prohibited.

Not for use on turf being grown for sale or other commercial use as sod.

Apply to moist or dry foliage. To avoid product pickup, lightly irrigate treated areas after application.

On short cut bentgrass (less than 3/4 inch) when temperatures are above 80°F, apply only to dry foliage.

Do not expose product to atmospheric moisture. For example, after application, fold down and seal top of the bag. This is particularly important under humid conditions.

Use only in clean, dry spreader. Do not leave product in spreader after application.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.

Do not apply this product when the wind direction is toward aquatic areas.

Do not apply this product using "belly grinder" or other handheld application equipment.

Do not make more than 6 applications per year.

Limitations, Restrictions, and Exceptions

FOR CONTROL

Apply at the DOUBLE RATE every 14-30 days until the disease is brought under control, then revert back to the preventative schedule. For more uniform particle distribution at the DOUBLE RATE, use a two-pass application in an alternate pattern at the NORMAL RATE.

To control severe brown patch infestations, use a non-fertilizer fungicide such as Andersons Golf Products Systemic Fungicide, Fungicide X, or Fungicide VII to limit the amount of fertilizer applied. This does not apply for the suppression of summer patch and red thread. Once the brown patch is controlled, prevent further infestations by following up with an Andersons Golf Products fungicide or combination fertilizer plus fungicide product.

Method

[Broadcast](#)

Rates

field_rates 0

-

Timings

Every 14-30 days until the disease is brought under control, then revert back to the preventive schedule.