

## **FOR ESTABLISHED TURF**

### General Information

#### DIRECTIONS FOR USE

It is in violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

#### WHEN TO APPLY

Apply prior to germination of weed grasses in the spring. Precise time of application for a specific area will vary due to climate and individual cultural practices.

Generally, application during February or March will give best results. Control can be expected by applications up to the time that Forsythias bloom in Spring. If in doubt, check with your Agricultural Extension Service. For newly seeded areas, apply as the last step before watering.

#### HOW TO APPLY

For best results, this product must be applied without skips or misses. Determine the correct amount for your lawn area. The best results are obtained by using the two-step procedure at the recommended half rate settings. Apply half the required amount in one direction and the other half at right angles to the first application. This procedure will minimize the chances of color streaking and erratic crabgrass control. Use the suggested spreader settings as a guide, being sure to check the application rate at the start of each application. At least 1/2 inch of water must be provided by rainfall or watering within 3 days of treatment in order to carry the chemical into the root zone of germinating weeds.

#### RESTRICTIONS

Do not enter or allow others to enter the treated area until dusts have settled.

If watering in is required after application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Do not apply this product in a way that will contact workers or other persons, either

directly or through drift. Only protected handlers may be in the area during application.

Aerial application is prohibited.

Golf Course Greens (band-treatments)

For band-treatments: Do not make more than 6 applications of siduron (of one product or combinations of products) to golf course greens per year.

## PRECAUTIONS

Non-target plants can be adversely affected by this product.

## Limitations, Restrictions, and Exceptions

The best results are obtained by using the two-step procedure at the recommended half rate settings. Apply half the required amount in one direction and the other half at right angles to the first application. This procedure will minimize the chances of color streaking and erratic crabgrass control. Use the suggested spreader settings as a guide, being sure to check the application rate at the start of each application. At least 1/2 inch of water must be provided by rainfall or watering within 3 days of treatment in order to carry the chemical into the root zone of germinating weeds.

Method

[N.A.](#)

Rates

[field rates 0](#)

•

Timings

[Apply prior to germination of weed grasses in the Spring. Precise time of application for a specific area will vary due to climate and individual cultural practices.](#)

[Generally, application during February or March will give best results. Control can](#)