

TREE CROPS

General Information

ProNatural zinc is manufactured and designed for foliar application on plants to prevent or correct zinc deficiencies that usually occur when plants are small and may limit crop growth and yields. ProNatural zinc is readily available and rapidly absorbed by the plant because of the natural complexing properties. It is designed for foliar, row or broadcast application and is water soluble and non-toxic to plants when applied as directed.

ProNatural ZINC is a yield-enhancing formulation that contains amino acid complexed zinc.

ProNatural ZINC is a proprietary formulation that has been specially reacted with emulsion polymerization that provides a wide range of mixing ease with other fertilizers or chemicals.

DIRECTIONS FOR USE

Dilute one part ProNatural ZINC with 20 or more parts water. Apply using spray method and in an adequate amount of water that will provide complete coverage of the plants. Use of a nonionic wetting agent may improve spray coverage of certain hard-to-wet plants. Do Not Apply Undiluted.

ProNatural ZINC can be included in a regular spray program on crops. Consult with a Wilbur-Ellis representative on compatibility with other spray materials. The rate of application will depend on the crop, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full-sized plants. Reduce the rates proportionately when spraying smaller plants.

ProNatural ZINC may be used on legumes, grain crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, vine crops, tropical and subtropical fruits, and many other crops.

Limitations, Restrictions, and Exceptions

The applications may be repeated at 2 to 4-week intervals through the growing season. On deciduous trees, 2 to 80 fluid ounces may be applied in the oil sprays at

the delayed dormant stage. Post-harvest application of 2 to 80 fluid ounces may be made while active, green leaves remain on the trees.

Method

[Foliar spray](#)

[Broadcast](#)

Rates

[field rates 0](#)

-

Timings

[After the beginning of active growth.](#)