

WARM SEASON GRASSES - CUDWEED, ETC.

General Information

FEATURES & USE TIPS:

- Combination fertilizer and Pendimethalin preemergent herbicide.
- Provides unsurpassed preemergent control of:

Annual Grassy Weeds:

Crabgrass

Foxtail

Fall panicum

Goosegrass

Barnyardgrass

Poa annua

Dicot Weeds (before they germinate):

Chickweed

Knotweed

Hop clover

Oxalis

Purshlane

Henbit

Prostrate Spurge

Cudweed

Eveningprimrose

- Consistently sized particles reduce segregation potential
- Small particle formulation provides more particles per square inch for better herbicide distribution.
- Nitrogen content includes 40% Poly-S, specially formulated for superior slow release nitrogen feeding.

Seeding or Sprigging intervals after treatment

Delay reseeding or winter overseeding for at least 3 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

Treatment interval after Seeding or Sprigging

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Refer to label for Spreader settings, application and instructions.

Limitations, Restrictions, and Exceptions

Apply a repeat application of 145-203 lbs/acre after 6-8 weeks if the lower rate was used initially.

Method

[Broadcast](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Timings

[Prior to weed germination.](#)