

MEDIUM APPLICATION RATE

General Information

DIRECTIONS FOR USE

Apply only as directed. Apply during the growing season.

FLORIDA APPLICATIONS:

We recommend that you follow the Florida Green Industries BMP's at

http://www.flaes.org/pdf/BMP_Book_final.pdf

We recommend that you follow the Florida Golf Course BMP's at

<http://www.flaes.org/pdf/glfbmp07.pdf>

For all applications:

- Apply to dry foliage.
- Water in after application with a minimum of 0.2 inches to help move particles into the turf canopy.
- Do not collect clippings for at least one mowing following application to avoid particle pick-up.

Cool Season Grasses:

- Apply at the LOW or MEDIUM RATE every 6-8 weeks during the period of active growth.
- Apply at the LIGHT RATE when temperatures exceed 85°F.

Warm Season Grasses:

- Apply at the LOW to HIGH RATE every 4-6 weeks during the period of active growth.

Notice: This product contains the micronutrient iron for extra greening and to help prevent iron deficiency. Iron may stain concrete and other surfaces. Misapplied product to these surfaces should be removed promptly by sweeping or blowing. Do not wash off with water.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk or street, back onto your lawn.

Limitations, Restrictions, and Exceptions

For all applications:

- For use on fairways and roughs.
- DO NOT USE ON GRENS
- Do not apply with a drop-type spreader.
- Do not apply at rates lower than recommended to prevent mottling.
- Apply to dry foliage.
- Water in after application with a minimum of 0.2 inches to help move particles in the turf canopy and to avoid possible staining.
- Do not collect clippings for at least one mowing following application to prevent particle pick-up.

Cool Season Grasses

- Apply at LOW or MEDIUM RATE.
- Apply at HIGH or HEAVY RATE in spring or fall deeper greening or when a higher potassium fertilization is desired.
- Exception: Apply at LOW RATE in mid-summer when temperatures are above 80 degrees F.

Warm Season Grasses

- Apply at LOW or MEDIUM RATE.
- HIGH or HEAVY RATE may be used in early spring to help bring grasses out of dormancy (such as bermudagrass) or when a higher potassium fertilization is desired.
- Exception: Apply at LOW RATE during mid-summer when temperatures are

extreme (90 degrees F or above).

Method

[Broadcast](#)

Rates

[field rates 0](#)

•

Timings

[N.A.](#)