

## **LEAF SPOT - TO PREVENT**

### General Information

### DIRECTIONS FOR USE

Use of this product at residential sites is prohibited.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

### Entry Restrictions

Do not enter or allow others to enter the treated area until dusts have settled.

This fungicide product prevents and controls the following turfgrass diseases:

Brown patch

Leaf spot (*Helminthosporium* spp.)

Red leaf spot

Pink snow mold / Fusarium patch (*Microdochium nivale*)

Gray snow mold (*Typhula* spp.)

For use on fairways, putting greens and other turf areas consisting of Kentucky bluegrass, bentgrass, perennial ryegrass, fine fescue and bermudagrass, or where these mixtures predominate. Product may also be applied to St. Augustinegrass turf areas.

### HOW TO USE

For LIGHT, NORMAL or HEAVY RATE applications (0.5, 1.0 or 1.5 oz ai/1000 sq ft), apply to moist or dry foliage for summer diseases. Applications need not be watered in. Apply to dry foliage for snow mold prevention/control.

Delay mowing for 24 hours after application to prevent particle pickup by mowers.

Expected time disease activity:

Spring: leaf spot, pink snow mold (*Fusarium* patch), gray snow mold

Summer: leaf spot and brown patch

Fall: leaf spot and pink snow mold (*Fusarium* patch)

Winter: pink snow mold (Fusarium patch), and gray snow mold

Allow one week between the application of this and other control products. This guideline may be relaxed where a severe disease attack requires immediate treatment.

#### NOTE TO USER

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, estuary, reservoir, river, permanent stream, marsh or natural pond, or commercial fish pond there must be at least a 25 foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.

Do not apply this product when the wind direction is toward aquatic areas.

May apply up to but no more than 24 pounds active ingredient (ai) annually. Do not apply more than 6 times per year.

Do not apply this product using a "belly grinder" or other handheld application equipment.

#### Limitations, Restrictions, and Exceptions

##### Leaf Spot

To prevent, apply to moist or dry foliage at the LIGHT RATE (0.5 oz. ai/1000 sq. ft.) when disease activity is anticipated. Repeat if needed after 3 to 4 weeks.

Light Rate: 2.4 lbs / 1000 sq ft

##### Method

[Broadcast](#)

##### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

Timings

[When disease activity is anticipated.](#)