

## **ESTABLISHED COOL AND WARM SEASON GRASSES - NORMAL RATE**

### General Information

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not apply this product in a way that will contact workers or other person, either directly or through drift. Only protected handlers may be in the area during application.

Controls crabgrass and goosegrass as they sprout.

This product may be only applied by a tractor-drawn spreader or by push type spreader.

For use only on golf courses (greens, tees and bentgrass fairways only).

Application on bentgrass fairways is limited to the following states: OH, PA, NY, MI, CT, MA, IN, IL, NJ, WV, MN, WI, VT, NH, RI, DE, MD, VA. Applications on bentgrass fairways may only be made in the Fall and are limited to one application per year.

Applications to greens and tees are limited to two per year. The maximum application rate is 458 lbs./acre or 10.5 lb./1,000 sq. ft. of product (6 lbs oxadiazon ai per acre) per year, except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 610 lbs./acre or 14 lb./1,000 sq. ft. of product (8 lbs oxadiazon ai per acre) per year. The maximum single application rate is 305 lbs./acre or 7 lbs./1,000 sq. ft. of product (4 lbs. oxadiazon ai per acre).

#### PRECAUTIONS

**IMPORTANT: THIS PRODUCT WILL NOT WORK UNLESS IT IS WATERED IN FOR 10-15 MINUTES. FOR SAFETY REASONS, WATER THIS PRODUCT IN AS SOON AS POSSIBLE FOLLOWING APPLICATION FOR 10-15 MINUTES AND DO NOT ALLOW CHILDREN OR**

PETS ON TREATED AREAS UNTIL DRY FOLLOWING THE WATERING.

WATERING-IN MAY BE PERFORMED BY THE COMMERCIAL APPLICATOR OR THE COMMERCIAL APPLICATOR MUST PROVIDE THE ABOVE WATERING-IN INSTRUCTIONS TO THE RESIDENT IN WRITING USING BOLD FACE TYPE AND/OR RED PRINT OR OTHER METHODS TO ENSURE WARNING IS PROMINENTLY DISPLAYED ON THE MATERIAL PROVIDED TO THE RESIDENT.

Do not use on home lawns, parks, recreation areas, or other public sites.

Limitations, Restrictions, and Exceptions

ESTABLISHED COOL AND WARM-SEASON GRASSES

This product may be used on bentgrass and bermudagrass putting greens only under conditions of very heavy goosegrass infestations and where the herbicidal side effects are tolerable.

This product may cause yellowing and/or thinning for up to 4 weeks on bentgrass and bermudagrass greens.

Do not use this product when the temperature is consistently below 50°F or above 80°F.

Not for use where *Poa annua*, annual ryegrass, or fine fescue are the desired turf or on dichondra or similar ground covers.

Complete any raking or aerifying of the turf before treatment so that the protective chemical barrier is not destroyed.

For best results apply to DRY FOLIAGE and ensure that the product granules reach the soil surface by removing debris and mowing if necessary prior to treatment.

Apply uniformly at the NORMAL RATE (1X) in the Spring (South: January to early April; North: March to early May) prior to crabgrass or goosegrass germination.

ADDITIONAL SEEDING, OVERSEEDING, SPRIGGING, AND STOLONIZING INFORMATION

Do not use on newly seeded bermudagrass until after the second mowing.

Aerification, coring or other practices which break the herbicidal barrier will increase germination if performed just before seeding.

Residual effectiveness requires a delay of 5 growing months after the last treatment before overseeding. On newly sprigged bermudagrass wait until the grass has filled in and stolons have rooted down before application.

Where large areas of bermudagrass have been severely thinned (25% or more), withhold application until desired fill-in and rooting in of the stolons are achieved.

When applying the product at NORMAL RATE (1X), a HALF WIDTH application can be made. The HALF WIDTH application will ensure a more uniform coverage and particle distribution by reducing the effect of spreader variation. If the HALF WIDTH application is chosen, be sure to use the HALF WIDTH setting under the NORMAL RATE (1X) as shown in the spreader table at end of label.

#### Method

[Broadcast](#)

#### Rates

[field rates 0](#)

[field rates 1](#)

- 

#### Timings

[Apply uniformly in the Spring \(South: January to early April; North March to early May\) prior to crabgrass or goosegrass germination.](#)