

## **REDUCE YELLOWING OR INJURY - HALF RATE**

### General Information

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not apply this product in a way that will contact workers or other person, either directly or through drift. Only protected handlers may be in the area during application.

Controls crabgrass and goosegrass as they sprout.

This product may be only applied by a tractor-drawn spreader or by push type spreader.

For use only on golf courses (greens, tees and bentgrass fairways only).

Application on bentgrass fairways is limited to the following states: OH, PA, NY, MI, CT, MA, IN, IL, NJ, WV, MN, WI, VT, NH, RI, DE, MD, VA. Applications on bentgrass fairways may only be made in the Fall and are limited to one application per year.

Applications to greens and tees are limited to two per year. The maximum application rate is 458 lbs./acre or 10.5 lb./1,000 sq. ft. of product (6 lbs oxadiazon ai per acre) per year, except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 610 lbs./acre or 14 lb./1,000 sq. ft. of product (8 lbs oxadiazon ai per acre) per year. The maximum single application rate is 305 lbs./acre or 7 lbs./1,000 sq. ft. of product (4 lbs. oxadiazon ai per acre).

## PRECAUTIONS

IMPORTANT: THIS PRODUCT WILL NOT WORK UNLESS IT IS WATERED IN FOR 10-15 MINUTES. FOR SAFETY REASONS, WATER THIS PRODUCT IN AS SOON AS POSSIBLE FOLLOWING APPLICATION FOR 10-15 MINUTES AND DO NOT ALLOW CHILDREN OR PETS ON TREATED AREAS UNTIL DRY FOLLOWING THE WATERING.

WATERING-IN MAY BE PERFORMED BY THE COMMERCIAL APPLICATOR OR THE COMMERCIAL APPLICATOR MUST PROVIDE THE ABOVE WATERING-IN INSTRUCTIONS TO THE RESIDENT IN WRITING USING BOLD FACE TYPE AND/OR RED PRINT OR OTHER METHODS TO ENSURE WARNING IS PROMINENTLY DISPLAYED ON THE MATERIAL PROVIDED TO THE RESIDENT.

Do not use on home lawns, parks, recreation areas, or other public sites.

### Limitations, Restrictions, and Exceptions

#### Reduce Yellowing or Injury

To reduce the potential for yellowing or injury on established bentgrass greens, bermudagrass greens, and perennial ryegrass winter overseedlings, apply an initial treatment to dry foliage at the HALF RATE (Vz X). Apply a repeat treatment 10-14 days later at the HALF RATE (Vz X).

Where herbicidal effects at the NORMAL RATE (1X) can be tolerated, this product may be applied at the HALF WIDTH to ensure proper particle coverage.

#### Method

[Broadcast](#)

#### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

#### Timings

[N.A.](#)