

FOR USE ON RESIDENTIAL AND ORNAMENTAL TURFGRASS SITES AND SOD FARMS (COOL-SEASON GRASSES OTHER THAN BENTGRASS)

General Information

BK 800 controls perennial broadleaf weeds and undesirable woody plants established in noncropland. It is effective for buckbrush, poison ivy, multiflora rose, and sumac established in the uncultivated areas presented below:

UNCULTIVATED AGRICULTURAL AREAS AND UNCULTIVATED NONAGRICULTURAL AREAS:

A. Noncropland Sites

- Barrier strips
- Farmyards
- Fencerows or fence lines
- Firebreaks
- Highway rights-of-way (principal, interstate, county, private, and unpaved roads): Roadsides, roadside ditches, road shoulders, road embankments, dividers, and medians.
- Industrial sites: Lumberyards, tank farms, fuel or equipment storage areas.
- Municipal, state, and federal lands: Airports and military installations
- Railroad rights-of-way
- Recreation areas: Fairgrounds, golf courses, parks, and areas adjacent to athletic fields.
- Utility rights-of-way: Telephone, pipeline, electrical powerlines, and communication transmission lines

B. Prohibitions for Noncropland Sites.

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (noncropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.

INSTRUCTIONS:

To effectively control brush, all leaves, stems and suckers should be thoroughly wetted to the ground. Apply when plants come into full leaf (spring) until plants begin to go dormant. Best results are obtained when brush and broadleaf weeds are young and actively growing. Do not cut brush until the herbicide has translocated throughout the plant causing root death.

USE PRECAUTIONS:

Do not apply this product through any type of irrigation system.

Limitations, Restrictions, and Exceptions

FOR USE ON RESIDENTIAL AND ORNAMENTAL TURFGRASS SITES AND SOD FARMS (COOL-SEASON GRASSES OTHER THAN BENTGRASS)

For Use on Residential and Ornamental Turfgrass Sites and Sod Farms (Cool-Season Grasses Other Than Bentgrass):

USE PRECAUTIONS:

Do not use on carpetgrass, dichondra, St. Augustine grass, bentgrass, nor on lawns or turf where desirable clovers are present. Use only lawn type sprayers. Do not exceed specified dosages for any area. Do not apply to newly seeded grasses until well established. Do not spray when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application.

Do not use this product on or near desirable plants, including within the drip line of desirable trees and shrubs, since injury may result.

INSTRUCTIONS:

Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Avoid spraying during long, excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours after application.

Apply 2.0 to 3.0 pints of product in 20 to 260 gallons of water per acre (0.75 to 1.1 fluid ounces of product in 0.5 to 6.0 gallons of water per 1,000 square feet). Use higher rates when using the higher volume of water per acre.

Limitations on broadcast treatments for ornamental turfgrass and sod farms:

The maximum application rate is 3.0 pints of product per acre per application. For ornamental turfgrass, the maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. For sod farms, the maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 6.0 pints of product per acre.

Controlled Droplet Applicator (CDA) – Add 1.5 pints to the Herbi container and fill with water. Spray contents over 33,000 square feet.

Avoid overlapping between spray patterns.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[Postemergence \(Weed\)](#)