

MICE

General Information

GENERAL INFORMATION

Kills rats and mice.

Kills warfarin-resistance norway rats.

Norway rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

USE RESTRICTIONS

For control of Norway rats, roof rats and house mice. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. When used in USDA-inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait. Do not use in sewers.

Urban Areas

This product may be used in and around the periphery of homes, industrial, commercial, and public buildings. May also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. May also be used in alleys.

Non-Urban Areas: This product may be used inside of homes and agricultural buildings

SELECTION OF TREATMENT AREAS

Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

Limitations, Restrictions, and Exceptions

MICE

Place at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

FOLLOW-UP

Replace contaminated or spoiled bait immediately. Using gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Method

[Bait](#)

Timings

[N.A.](#)