

NON-CROP SITES - WEEDS CONTROL (1/2 TO 1 OUNCE PER ACRE)

General Information

IMPORTANT PRECAUTIONS

DO NOT USE ON FOOD OR FEED CROPS EXCEPT AS RECOMMENDED BY THE LABEL. Injury to or loss of desirable trees or other plants may result if the precautions listed below are not followed.

- Do not apply AmTide MSM 60 DF Herbicide (except as recommended), or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the product may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas except as recommended by the label.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water.
- Keep from contact with fertilizers, insecticides, fungicides and seeds.

Spraying and mixing equipment used with AmTide MSM 60 DF Herbicide must not be used for subsequent applications to food or feed crops with the exception of pastures, rangeland, and wheat, as low rates of AmTide MSM 60 DF HERBICIDE can kill or severely injure most food or feed crops.

AmTide MSM 60 DF should be used only in accordance with recommendations on the label. AmTide, LLC will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by AmTide, LLC. User assumes all risks associated with such non-recommended use.

TANK MIXES

AmTide MSM 60 DF HERBICIDE may be tank mixed with other Herbicides registered for the use sites described in the label. Use only those tank mix partners which are labeled for the appropriate use site. When tank mixing, use the most restrictive

label limitations for each of the products being used in the tank mix.

IMPORTANT INFORMATION

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.
- Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

AmTide MSM 60 DF HERBICIDE is recommended for use on land primarily dedicated to the production of wheat, barley, fallow, pasture and rangeland.

AmTide MSM 60 DF HERBICIDE is recommended for use on wheat, barley, fallow, pasture, and rangeland in most states. Check your state extension or Dept. of Agriculture before use to be certain AmTide MSM 60 DF HERBICIDE is registered in your state. AmTide MSM 60 DF HERBICIDE is not registered for use in Alamosa, Conejos, Costilla, Rio Grande and Saquache counties of Colorado.

AmTide MSM 60 DF HERBICIDE is a dry-flowable granule that controls weeds in wheat (including durum), barley, fallow, pasture, and rangeland grasses. AmTide MSM 60 DF HERBICIDE is mixed in water or can be preslurried in water and added to

liquid nitrogen carrier solutions and applied as a uniform spray mix unless otherwise specified on the label. AmTide MSM 60 DF HERBICIDE is non-corrosive, nonflammable, nonvolatile, and does not freeze.

AmTide MSM 60 DF HERBICIDE controls weeds by postemergence activity. For best results, apply AmTide MSM 60 DF HERBICIDE to young, actively growing weeds. The use rates depend upon the weed spectrum and size of weeds at application. The degree and duration of control may depend on the following factors:

- Weed spectrum and infestation intensity
- Weed size at application
- Environmental condition at and following treatment

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

AmTide MSM 60 DF is absorbed through the foliage of broadleaf weeds, rapidly inhibiting their growth. Leaves of susceptible plants appear chlorotic from 1 to 3 weeks after application and the growing point subsequently dies.

Application of AmTide MSM 60 DF provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

AmTide MSM 60 DF may injure crops that are stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, different varieties of the crop may be sensitive to treatment with AmTide MSM 60 DF under otherwise normal conditions. Treatment of such varieties may injure crops.

In warm, moist conditions, the expression of Herbicide symptoms is accelerated in weeds; in cold, dry conditions, expression of Herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to AmTide MSM 60 DF.

Weed control may be reduced if rainfall or snowfall occurs soon after application.

Limitations, Restrictions, and Exceptions

NON-CROP SITES

Application Information

AmTide MSM 60 DF HERBICIDE is recommended for general weed control on private, public and military lands as follows: Uncultivated areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas, etc.); uncultivated agricultural areas - non-crop producing (such as farmyards, fuel storage areas, fence rows, soil bank land, barrier strips, etc.); industrial sites - outdoor (such as lumberyards, pipeline and tank farms, etc.). It is also recommended for the control of certain noxious and troublesome weeds.

AmTide MSM 60 DF HERBICIDE may be applied in tank mixture with other Herbicides labeled for use on non-crop sites. Fully read the labels and follow all the directions and restrictions on each label.

Applications may be made by ground or air. Use a sufficient volume of water to ensure thorough coverage of the target vegetation with the application equipment being used.

Application Timing

For best results, AmTide MSM 60 DF HERBICIDE should be applied postemergence to young, actively growing weeds. Applications may be made at any time of the year, except when the ground is frozen.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[Postemergence \(Weed\)](#)